


Bacteria never swim alone

July 14 2017



Many animal species display flocking behaviour, but the fact that microorganisms do is not as well known. Researchers at Lund University in Sweden have now shown that algae and bacteria form flocks at very low concentrations of individuals, a finding that could increase our future understanding of how the organisms infect their host animals.

Flocking behaviour in [animals](#) seemingly arises spontaneously in a group

of independent [individuals](#) without a clear leader. This behaviour occurs among all types of organisms, from bacteria to people. One hypothesis, therefore, is that there are fundamental principles for flock building that are not dependent on single individuals.

Researchers at Lund University, in cooperation with colleagues from the UK and France, have now found that flocking behaviour among microorganisms is more advanced than we previously thought.

"Our research is a physical explanatory model of how microorganisms move. From a biological perspective, it is useful to examine the evolutionary basis for flocking behaviour among bacteria, as the connections can increase our understanding of the course of infectious diseases," says Joakim Stenhammar, chemistry researcher at Lund University.

When a person or animal swims, they create backwashes or wakes that others can sense. The researchers have now created a theoretical model that describes how single microorganisms communicate with each other via the backwashes that each organism creates. The physical principle differs from ordinary backwashes, but these flows enable the bacteria to sense each other's presence and affect each other at very low concentrations. In the light of this, microorganisms cannot be described as isolated individuals.

It was previously known that certain swimming bacteria, such as *E. coli* and *Salmonella*, form flocks at high concentrations. In the new study, Stenhammar and his colleagues have shown that it is only at extremely low concentrations – less than ten per cent of what was previously thought – that [bacteria](#) can be considered as individuals.

"In contrast to an individual bacterium, flocks can move in a synchronised way over long length scales and several times faster than a

single bacterium," says Joakim Stenhammar.

"Our research adds another piece of the puzzle to our understanding of how flocking [behaviour](#) works in biological systems, and the model can be applied to a large number of swimming [microorganisms](#)," says Joakim Stenhammar.

More information: Joakim Stenhammar et al. Role of Correlations in the Collective Behavior of Microswimmer Suspensions, *Physical Review Letters* (2017). [DOI: 10.1103/PhysRevLett.119.028005](https://doi.org/10.1103/PhysRevLett.119.028005)

Provided by Lund University

Citation: Bacteria never swim alone (2017, July 14) retrieved 18 April 2024 from <https://phys.org/news/2017-07-bacteria.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.