


# A cosmic barbecue: Researchers spot 60 new 'hot Jupiter' candidates

July 6 2017

---



A hot Jupiter at various phases of its orbit. The sizes of the star and planet and the separations between them are to scale for a typical hot Jupiter. The amount of reflected starlight that is observed depends on the planet's position within its orbit and the inclination of the orbit with respect to the observer. Credit: Millholland/Yale University

Yale researchers have identified 60 potential new "hot Jupiters"—highly irradiated worlds that glow like coals on a barbecue grill and are found

orbiting only 1% of Sun-like stars.

Hot Jupiters constitute a class of [gas giant planets](#) located so close to their parent [stars](#) that they take less than a week to complete an orbit. Second-year Ph.D. student Sarah Millholland and astronomy professor Greg Laughlin identified the [planet candidates](#) via a novel application of big data techniques. They used a supervised machine learning algorithm—a sophisticated program that can be trained to recognize patterns in data and make predictions—to detect the tiny amplitude variations in observed [light](#) that result as an orbiting planet reflects rays of light from its host star.

"Sarah's work has given us what amounts to a 'class portrait' of extrasolar planets at their most alien," said Laughlin. "It's amazing how the latest techniques in machine learning, compounded with high-performance computing, are allowing us to mine classic data sets for extraordinary discoveries."

Millholland recently presented the research at a Kepler Science Conference at the NASA Ames Research Center in California. She and Laughlin are authors of a study about the research, which has been accepted for publication in the *Astronomical Journal*.

Millholland and Laughlin searched systematically for reflected [light signals](#) in the observations of more than 140,000 stars from four years of data from NASA's Kepler mission. The Kepler spacecraft is best known for enabling the detection of thousands of exoplanets that transit their host stars. During a transit, a planet passes in front of a star and causes a periodic dip in the observed starlight.



The graphic shows the reflected light signals for the 60 candidate planets. The colorations of the stars are appropriate to their temperatures, and the sizes of the stars and candidate planets are to scale. Credit: Yale University

Reflected light signals can be difficult to distinguish from stellar or instrumental variability, the researchers said, but a big data approach enabled them to pull out the faint signals. They generated thousands of

synthetic datasets and trained an algorithm to recognize the properties of the reflected light signals in comparison to those with other types of variability.

The Yale technique pioneers a new discovery method that identifies more planets from the publicly available Kepler data, said the researchers. "I've been told by members of the Kepler science team that a search for reflected star-shine was part of the early renditions of the Kepler pipeline," Millholland said. "They called it the Reflected Light Search, or RLS module. In this sense, we're simply addressing one of the original intentions for the Kepler data."

The reflected light signals hold rich information about the [planets'](#) atmospheres, according to the researchers. They contain characteristics such as cloud existence, atmospheric composition, wind patterns, and day-night temperature contrasts.

The researchers also note that the 60 planet candidates will require follow-up observations for confirmation. This will come in the form of Doppler velocity measurements.

The Doppler velocity method is a well-established technique that enables the detection of wobbling motion in a star due to the gravitational influence of an orbiting planet. Since hot Jupiters are so massive and close to their stars, the stellar wobbles they induce are large and readily detectable.

A new, Yale-designed instrument known as EXPRES, which is being installed on the Discovery Channel Telescope in Arizona, may attempt to make confirmations later this year.

A 3D manipulable diagram of a hot Jupiter in various phases of its orbit is available at: [smilholland.github.io/Phase\\_Curve\\_Demo/](https://smilholland.github.io/Phase_Curve_Demo/)

**More information:** Supervised Learning Detection of Sixty Non-Transiting Hot Jupiter Candidates, [arxiv.org/abs/1706.06602](https://arxiv.org/abs/1706.06602)

Provided by Yale University

Citation: A cosmic barbecue: Researchers spot 60 new 'hot Jupiter' candidates (2017, July 6)  
retrieved 23 April 2024 from  
<https://phys.org/news/2017-07-cosmic-barbecue-hot-jupiter-candidates.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--