

High-speed FM-AFM and simulation reveal atomistic dissolution processes of calcite in water

July 27 2017



(a) Atomistic model of calcite surface. (b) The dissolution processes of calcite surface in water observed with high-speed FM-AFM. It is observed that the step is moving from lower-right to upper-left. Along the step is also seen the transition region. (c) Averaged height profile measured along the line PQ indicated in (b). The height of a monolayer step is ~0.3 nm, but that of the transition region is smaller. A terrace described in the Figure indicates a flat area at the atomic level on the crystal surface. The upper terrace is higher by one



monolayer of CaCO3 than the lower terrace. Credit: Kanazawa University

Calcite is one of the most abundant components of the Earth's crust, constituting the largest carbon reservoir in the global carbon cycle. Thus, large-scale dissolution of calcite would have enormous impact on the weather, geography and aquatic environment, for example, changes in the carbon dioxide concentration of the air and the acidity of the ocean. The dissolution mechanism of calcite has importance in geologic carbon sequestration (GCS) technology to capture carbon dioxide from the air and to store it underground. In order to precisely predict such a large-scale and long-term phenomenon, the dissolution mechanism of calcite should be understood at an atomic level in a precise manner.

When a crystal of calcite is immersed in water (Figure 1a), it is observed that a monolayer of ~0.3 nm thickness is formed on the surface exposed to water—this is called the step edge. The crystal dissolution proceeds as desorption of atoms from the step edge to aqueous solution. Therefore, understanding of atomistic events at step edges is essential for elucidation of the dissolution processes. Nonetheless, due to the limitations of measurement technologies, it was difficult to observe highspeed structural changes associated with the atomistic dissolution process . Thus, many aspects of crystal growth and dissolution mechanisms, including those of calcite, remained unclear.

Atomic force microscopy (AFM) is capable of observing the surface morphology of insulating materials. Therefore, AFM is thought to be a measurement technique that may have great potential to solve the problem described above. Nonetheless, conventional AFMs do not have enough spatial or temporal resolution for this purpose.





With a model that places a monolayer of Ca(OH)2 in proximity of a step at the boundary of upper terrace and lower terrace, molecular dynamics simulation was performed for about 7.5 ns to confirm that the monolayer of Ca(OH)2 existed stably adjacent to the step without being separated from the crystal surface. Credit: Kanazawa University

Researchers of Kanazawa University, Japan, have led the development of technologies for frequency modulation AFM (FM-AFM) over the years, and have advanced the temporal resolution to ~1 s/frame from the current standard of ~1 min/frame. The international research team succeeded in the direct observation of the dissolution processes of the calcite surface in water as well as of structural changes around step edges at the atomistic level for the first time. Moreover, from the FM-AFM images, the team has found that the transition region of a few nanometers' width along a step is formed as an intermediate state in the dissolution processes (Figure 1b). The formation of this transition region was not foreseen by previous studies, and without the high-speed FM-AFM, it would not have been discovered. In addition, in order to elucidate the origin of the transition region and dissolution mechanism, the team examined the validity of various transition region models by density functional theory calculations and by molecular dynamics simulations (Figure 2). It was found that the transition region would most



likely be a Ca(OH)2 monolayer formed as an intermediate state in the dissolution processes of calcite. Based on these results, the team proposes a dissolution mechanism at atomistic level as follows (Figure 3).

- 1. At the step edges, dissociative adsorption of a water molecule leads to generation of ion pair of surface-bound CaOH+ and free HCO3-.
- 2. HCO3- is decomposed and surface-bound Ca(OH)2 and free CO2 are formed.
- 3. Repetition of these reactions forms the transition region consisting of a Ca(OH)2 monolayer at the step edge.
- 4. At the peripheries of the transition region, stability of the surfaceadsorbed Ca(OH)2 molecules depends on the distance from the step edge, and at a certain distance (typically, a few nanometers), Ca(OH)2 dissociates.

To the team's knowledge, this is the very first proposal for the dissolution processes at atomistic level based on such direct experimental evidence. Moreover, this is also the first proposal for the dissolution mechanism of calcite with the formation of the transition region taken into consideration. Thus, the team believes that the present study promotes the understanding of calcite dissolution mechanism at atomistic level to a great extent.





Atomistic dissolution model of calcite surface in water. Credit: Kanazawa University

The precise understanding of the dissolution processes of calcite at an atomistic level may enable researchers to understand the physical meanings of empirical parameters used for simulations of the dissolution processes at a macroscopic level. This may also lead to accurate prediction of dissolution behaviors in various solution environments in nature, and the present study is expected to make contributions to better prediction accuracy of the global carbon cycle. Furthermore, the high-speed FM-AFM developed and reported in this study will be applicable not only to studies of the dissolution processes of calcite but to those of crystal growth, dissolution and self-assembly of a variety of minerals and organic as well as biological molecules. It will also be quite useful for observation and investigation of a wide variety of solid-liquid interface phenomena at an atomistic level such as metal corrosion, catalytic reaction, etc. Since there were no appropriate direct observation means available for those phenomena, the present high-speed FM-AFM is



expected to pave the way for discoveries of various phenomena so far unknown.

More information: Kazuki Miyata et al, Dissolution Processes at Step Edges of Calcite in Water Investigated by High-Speed Frequency Modulation Atomic Force Microscopy and Simulation, *Nano Letters* (2017). DOI: 10.1021/acs.nanolett.7b00757

Provided by Kanazawa University

Citation: High-speed FM-AFM and simulation reveal atomistic dissolution processes of calcite in water (2017, July 27) retrieved 6 May 2024 from <u>https://phys.org/news/2017-07-high-speed-fm-afm-simulation-reveal-atomistic.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.