

Neutrons detect elusive Higgs amplitude mode in quantum material

July 5 2017, by Sara Shoemaker



The ORNL-led research team selected a crystal composed of copper bromide -- because the copper ion is ideal for studying exotic quantum effects -- to observe the elusive Higgs amplitude mode in two dimensions. The sample was examined using cold neutron triple-axis spectrometer beams for neutron scattering at the High Flux Isotope Reactor. Credit: Genevieve Martin, Oak Ridge National Laboratory/Dept. of Energy


A team led by the Department of Energy's Oak Ridge National Laboratory has used sophisticated neutron scattering techniques to detect an elusive quantum state known as the Higgs amplitude mode in a two-dimensional material.

The Higgs [amplitude](#) mode is a condensed matter cousin of the Higgs boson, the storied quantum particle theorized in the 1960s and proven experimentally in 2012. It is one of a number of quirky, collective modes of matter found in materials at the quantum level. By studying these modes, condensed matter researchers have recently uncovered new quantum states known as quasiparticles, including the Higgs mode.

These studies provide unique opportunities to explore quantum physics and apply its exotic effects in advanced technologies such as spin-based electronics, or spintronics, and quantum computing.

"To excite a material's quantum quasiparticles in a way that allows us to observe the Higgs amplitude mode is quite challenging," said Tao Hong, an instrument scientist with ORNL's Quantum Condensed Matter Division.

Although the Higgs amplitude mode has been observed in various systems, "the Higgs mode would often become unstable and decay, shortening the opportunity to characterize it before losing sight of it," Hong said.



ORNL's Tao Hong analyzed a copper bromide compound's low-energy behavior during a neutron scattering experiment at the lab's High Flux Isotope Reactor that yielded the elusive Higgs amplitude mode in two dimensions with no decay. Credit: Genevieve Martin, Oak Ridge National Laboratory/Dept. of Energy


The ORNL-led team offered an alternative method. The researchers selected a crystal composed of copper bromide, because the copper ion is ideal for studying exotic quantum effects, Hong explained. They began the delicate task of "freezing" the material's agitating quantum-level particles by lowering its temperature to 1.4 Kelvin, which is about minus 457.15 degrees Fahrenheit.

The researchers fine-tuned the experiment until the particles reached the phase located near the desired [quantum critical point](#)—the sweet spot

where collective quantum effects spread across wide distances in the material, which creates the best conditions to observe a Higgs amplitude mode without decay.

With neutron scattering performed at ORNL's High Flux Isotope Reactor, the research team observed the Higgs mode with an infinite lifetime: no decay.

"There's an ongoing debate in physics about the stability of these very delicate Higgs modes," said Alan Tennant, chief scientist of ORNL's Neutron Sciences Directorate. "This experiment is really hard to do, especially in a two-dimensional system. And, yet, here's a clear observation, and it's stabilized."



During the neutron scattering experiment, the sample containing copper ions exhibited exotic quantum properties as certain quasiparticles spin in a wave-like configuration, eventually revealing the Higgs amplitude mode. Credit: Oak Ridge National Laboratory/Dept. of Energy

The research team's observation provides new insights into the fundamental theories underlying exotic materials including superconductors, charge-density wave systems, ultracold bosonic systems and antiferromagnets.

"These breakthroughs are having widespread impact on our understanding of materials' behavior at the atomic scale," Hong added.

The study, titled, "Direct observation of the Higgs amplitude mode in a two-dimensional quantum antiferromagnet near the [quantum](#) critical point," was published in *Nature Physics*.

More information: Tao Hong et al, Higgs amplitude mode in a two-dimensional quantum antiferromagnet near the quantum critical point, *Nature Physics* (2017). [DOI: 10.1038/nphys4182](https://doi.org/10.1038/nphys4182)

Provided by Oak Ridge National Laboratory

Citation: Neutrons detect elusive Higgs amplitude mode in quantum material (2017, July 5) retrieved 28 January 2023 from <https://phys.org/news/2017-07-neutrons-elusive-higgs-amplitude-mode.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.