

Thawing permafrost releases old greenhouse gas

July 19 2017



View out of the window of the research aircraft Polar 5 during the flight campaign over the Mackenzie Delta. Credit: Picture: T. Sachs, GFZ

The thawing permafrost soils in the Arctic regions might contribute to the greenhouse effect in two respects: On the one hand, rising



temperatures lead to higher microbial methane production close to the surface. On the other hand, the thawing subsurface opens pathways for old, geologic methane. This is shown in a study in the Mackenzie Delta (Canada), conducted by scientists from the German Research Centre for Geosciences GFZ, the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research (AWI) and partners in the U.S. The study is published in the journal *Scientific Reports*.

Arctic permafrost acts like a gigantic cap of frozen material over mineral resources and fossil fuels. Scientists apprehend that thawing permafrost could lead to rising <u>methane emissions</u>. "We wanted to find out how much methane is released in a region and were looking for spatial patterns in gas emissions," says lead author Katrin Kohnert from GFZ's section for Remote Sensing. In order to do so, a team led by GFZ scientist Torsten Sachs performed a study in a 10,000 square kilometers wide area in Northern Canada. Using the research aircraft Polar 5 of AWI, the scientists conducted airborne measurements of gas concentrations in the air during two extensive flight campaigns in summer 2012 and 2013, respectively.

The result was a high-resolution (100 m x 100 m) methane flux map of the Mackenzie Delta. "We found strong emissions solely where the permafrost is discontinuous, meaning parts where the permafrost contains areas that are thawed permanently," says Katrin Kohnert. "We think that the methane comes predominantly from deeper geologic sources and not from recent microbial activity close to the surface." Even though the hotspots only occur on about 1 percent of the area, they contribute 17 percent to the annual methane emission estimate of the study area.

The conclusion of the authors: The warming climate triggers not only the natural production of biogenic <u>methane</u>, it can also lead to stronger emissions of fossil gas. This contributes significantly to the permafrost-



carbon-climate feedback. Kohnert: "Therefore, permafrost areas vulnerable to thawing warrant much more attention."



Map of the methane fluxes in the northern part of the study area. Credit: B. Juhls, GFZ



More information: Katrin Kohnert et al, Strong geologic methane emissions from discontinuous terrestrial permafrost in the Mackenzie Delta, Canada, *Scientific Reports* (2017). <u>DOI:</u> <u>10.1038/s41598-017-05783-2</u>

Provided by Helmholtz Association of German Research Centres

Citation: Thawing permafrost releases old greenhouse gas (2017, July 19) retrieved 26 April 2024 from <u>https://phys.org/news/2017-07-permafrost-greenhouse-gas.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.