


Quantum probes dramatically improve detection of nuclear spins

July 3 2017



A nitrogen-vacancy (dark blue) quantum probe in diamond (light grey) performing nanoscale nuclear magnetic resonance (NMR) on molecular hydrogen sitting on the diamond's surface. A green laser controls the quantum state of the probe, which is tuned to the resonant frequency of target nuclear spins. The probe responds to the nuclear spins of the hydrogen atoms and provides a direct measurement via the red light emitted. Credit: David A. Broadway/cqc2t.org

Researchers at the University of Melbourne have demonstrated a way to detect nuclear spins in molecules non-invasively, providing a new tool for biotechnology and materials science.

Important research in medicine and biology relies on nuclear magnetic resonance (NMR) spectroscopy, but until now, it has been limited in spatial resolution and typically requires powerful microwave fields. A team led by Professor Lloyd Hollenberg at the University of Melbourne has used a quantum probe to perform microwave-free NMR at the nanoscale. The results were published today in *Nature Communications*.

"This quantum probe delivers a dramatic improvement in NMR technology. In addition to being able to detect NMR in far smaller samples than conventional machines, our technique does not require the application of microwave fields that might disrupt biological samples" said Hollenberg, who is Deputy Director of the Centre for Quantum Computation and Communication Technology (CQC2T) and Thomas Baker Chair at the University of Melbourne.

"In NMR the goal is to detect the magnetic signal from the nuclei of the atoms comprising molecules. But the signal from the nuclear "spin" is very weak and conventional NMR machines require many millions of [nuclear spins](#) to detect anything. However, using the quantum properties of a 'defect' in diamond, our technique can detect much smaller volumes down to only thousands of spins."

The discovery may overcome significant limitations with conventional NMR methods, which depend on machines that can exceed 10 tonnes.

"The problem with the large NMR [machines](#) in widespread use today is that the signals we're trying to detect are extremely small, and the distance from the measurement device to the object being measured is very large," said Dr. Alastair Stacey, a CQC2T postdoctoral researcher.

"This creates two problems: The machine can only see a larger collection of molecules, reducing the accuracy of the measurement. It also has to use very strong microwaves and magnetic fields to reach the sample, but these processes are invasive and can affect delicate bio-samples, just like the microwave in your kitchen, particularly when trying to see the molecular structure of liquids."

Lead author James Wood describes the technique as "a dramatic simplification of the nuclear detection process, where we essentially shine light on an atomic-sized defect in diamond and observe its natural response, at a fundamentally quantum level, to the target nuclear spins nearby".

"A great benefit of our approach is that we don't interfere with the sample when imaging it."

The technique offers new opportunities for researchers.

"With these advances in [quantum](#) sensing technology, we are opening the door to a new world of scientific investigation that could lead us to gain a better understanding of the smallest building blocks of life," said Hollenberg.

More information: *Nature Communications* (2017). [DOI: 10.1038/NCOMMS15950](https://doi.org/10.1038/NCOMMS15950)

Provided by Centre for Quantum Computation & Communication Technology

Citation: Quantum probes dramatically improve detection of nuclear spins (2017, July 3) retrieved 5 May 2024 from <https://phys.org/news/2017-07-quantum-probes-nuclear.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.