


# Scientists produce robust catalyst to split water into hydrogen, oxygen

July 26 2017



A catalyst developed by Rice University and the University of Houston splits water into hydrogen and oxygen without the need for expensive metals like

platinum. This electron microscope image shows nickel foam coated with graphene and then the catalytic surface of iron, manganese and phosphorus. Credit: Desmond Schipper/Rice University

Splitting water into hydrogen and oxygen to produce clean energy can be simplified with a single catalyst developed by scientists at Rice University and the University of Houston.

The electrolytic film produced at Rice and tested at Houston is a three-layer structure of nickel, graphene and a compound of iron, manganese and phosphorus. The foamy nickel gives the film a large surface, the conductive graphene protects the nickel from degrading and the metal phosphide carries out the reaction.

The robust material is the subject of a paper in *Nano Energy*.


Rice chemist Kenton Whitmire and Houston electrical and computer engineer Jiming Bao and their labs developed the film to overcome barriers that usually make a catalyst good for producing either [oxygen](#) or hydrogen, but not both simultaneously.

"Regular metals sometimes oxidize during catalysis," Whitmire said.

"Normally, a hydrogen evolution reaction is done in acid and an oxygen evolution reaction is done in base. We have one material that is stable whether it's in an acidic or basic solution."

The discovery builds upon the researchers' creation of a simple oxygen-evolution catalyst revealed earlier this year. In that work, the team grew a catalyst directly on a semiconducting nanorod array that turned sunlight into energy for [solar water splitting](#).

Electrocatalysis requires two catalysts, a cathode and an anode. When placed in water and charged, hydrogen will form at one electrode and oxygen at the other, and these gases are captured. But the process generally requires costly metals to operate as efficiently as the Rice team's catalyst.



A film of high-surface-area nickel foam coated with graphene and a compound of iron, manganese and phosphorus serve as a water-splitting catalyst that can produce hydrogen and oxygen simultaneously. The material was created at Rice University and tested at the University of Houston. Credit: Jeff Fitlow/Rice University

"The standard for hydrogen evolution is platinum," Whitmire said.

"We're using Earth-abundant materials—iron, manganese and phosphorus—as opposed to noble metals that are much more expensive."

The new [catalyst](#) also requires less energy, Whitmire said. "If you want to make hydrogen and oxygen, you have to put in energy, and the more you put in, the less commercially viable it is," he said. "You want to do it at the minimum amount of energy possible. That's a benefit of our material: The overpotential (the amount of energy required to trigger electrocatalysis) is small, and quite competitive with other materials. The lower you can get it, the closer you come to making it as efficient as possible for water splitting."

Graphene, the atom-thick form of carbon, is key to protecting the underlying nickel. One to three layers of graphene are formed on the nickel foam in a chemical vapor deposition (CVD) furnace, and the iron, manganese and phosphorus are added on top of that, also via CVD and from a single precursor.

Tests by Bao's lab compared nickel foam and the phosphide both with and without graphene in the middle and found the conductive graphene lowered charge-transfer resistance for both hydrogen and oxygen reactions.

"Nickel is one of the best catalysts to make [graphene](#)," said co-lead author Desmond Schipper, a Rice graduate student. "Essentially, we're using the nickel to help improve the [nickel](#)." He said the manganese adds a level of protection as well.

Whitmire said the material is scalable and should find use in industries that produce [hydrogen](#) and oxygen or by solar- and wind-powered facilities that can use electrocatalysis to store off-peak [energy](#).

It may also be adapted to produce other advanced materials. "Our



method could be widely applicable to a large number of metal phosphide materials for catalysts—not just for [water splitting](#), but for a range of things," he said.

"A critical factor is that we're able to make phase-pure [materials](#) with different compositions. Currently, people have very little control over the phase they get on a surface, and in many cases they get a mixture. When that happens, they don't know which phase is actually responsible for the catalysis. With our process, they can know."

**More information:** Zhenhuan Zhao et al, Bifunctional metal phosphide FeMnP films from single source metal organic chemical vapor deposition for efficient overall water splitting, *Nano Energy* (2017). [DOI: 10.1016/j.nanoen.2017.07.027](https://doi.org/10.1016/j.nanoen.2017.07.027)

Provided by Rice University

Citation: Scientists produce robust catalyst to split water into hydrogen, oxygen (2017, July 26) retrieved 13 March 2024 from <https://phys.org/news/2017-07-scientists-robust-catalyst-hydrogen-oxygen.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--