

Scientists get best measure of star-forming material in galaxy clusters in early universe

July 20 2017



The Tadpole Galaxy is a disrupted spiral galaxy showing streams of gas stripped by gravitational interaction with another galaxy. Molecular gas is the required ingredient to form stars in galaxies in the early universe. Credit: Hubble Legacy Archive, ESA, NASA and Bill Snyder.

The international Spitzer Adaptation of the Red-sequence Cluster Survey (SpARCS) collaboration based at the University of California, Riverside has combined observations from several of the world's most powerful telescopes to carry out one of the largest studies yet of molecular gas - the raw material which fuels star formation throughout the universe - in three of the most distant clusters of galaxies ever found, detected as they appeared when the universe was only four billion years old.

Results were recently published in *The Astrophysical Journal Letters*. Allison Noble, a postdoctoral researcher at the Massachusetts Institute of Technology, led this newest research from the SpARCS collaboration.

Clusters are rare regions of the universe consisting of tight groups of hundreds of [galaxies](#) containing trillions of [stars](#), as well as hot gas and mysterious dark matter. First, the research team used spectroscopic observations from the W. M. Keck Observatory on Mauna Kea, Hawai'i, and the Very Large Telescope in Chile that confirmed 11 galaxies were star-forming members of the three massive clusters. Next, the researchers took images through multiple filters from NASA's Hubble Space Telescope, which revealed a surprising diversity in the galaxies' appearance, with some galaxies having already formed large disks with spiral arms.

One of the telescopes the SpARCS scientists used is the extremely sensitive Atacama Large Millimeter Array (ALMA) [telescope](#) capable of directly detecting radio waves emitted from the molecular gas found in galaxies in the [early universe](#). ALMA observations allowed the scientists to determine the amount of molecular gas in each galaxy, and provided the best measurement yet of how much fuel was available to form stars.

The researchers compared the properties of galaxies in these clusters

with the properties of "field galaxies" (galaxies found in more typical environments with fewer close neighbors). To their surprise, they discovered that [cluster galaxies](#) had higher amounts of molecular gas relative to the amount of stars in the galaxy, compared to field galaxies. The finding puzzled the team because it has long been known that when a galaxy falls into a [cluster](#), interactions with other cluster galaxies and hot gas accelerate the shut off of its star formation relative to that of a similar field galaxy (the process is known as *environmental quenching*).

"This is definitely an intriguing result," said Gillian Wilson, a professor of physics and astronomy at UC Riverside and the leader of the SpARCS collaboration. "If cluster galaxies have more fuel available to them, you might expect them to be forming more stars than [field galaxies](#), and yet they are not."

Noble, a SpARCS collaborator and the study's leader, suggests several possible explanations: It is possible that something about being in the hot, harsh cluster environment surrounded by many neighboring galaxies perturbs the molecular gas in cluster galaxies such that a smaller fraction of that gas actively forms stars. Alternatively, it is possible that an environmental process, such as increased merging activity in cluster galaxies, results in the observed differences between the cluster and field galaxy populations.

"While the current study does not answer the question of which physical process is primarily responsible for causing the higher amounts of molecular gas, it provides the most accurate measurement yet of how much [molecular gas](#) exists in galaxies in clusters in the early universe," Wilson said.

The SpARCS team has developed new techniques using infrared observations from NASA's Spitzer Space Telescope to identify hundreds of previously undiscovered clusters of galaxies in the early universe. In

the future, they plan to study a larger sample of clusters. The team has recently been awarded additional time on ALMA, the W. M. Keck Observatory, and the Hubble Space Telescope to continue investigating how the neighborhood in which a galaxy lives determines for how long it can form stars.

More information: A. G. Noble et al. ALMA Observations of Gas-rich Galaxies in $z \sim 1.6$ Galaxy Clusters: Evidence for Higher Gas Fractions in High-density Environments, *The Astrophysical Journal* (2017). [DOI: 10.3847/2041-8213/aa77f3](https://doi.org/10.3847/2041-8213/aa77f3)

Provided by University of California - Riverside

Citation: Scientists get best measure of star-forming material in galaxy clusters in early universe (2017, July 20) retrieved 24 April 2024 from <https://phys.org/news/2017-07-scientists-star-forming-material-galaxy-clusters.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
