

Possible first sighting of an exomoon

July 28 2017, by Bob Yirka



This artist's concept depicts a planetary system. Credit: NASA/JPL-Caltech

(Phys.org)—A team led by David Kipping of Columbia University has spotted what might be the first evidence of an exomoon. They have written a paper describing their findings and have uploaded it to the *arXiv* preprint server.

An exomoon is a moon orbiting an exoplanet. Many moons have been

found in our solar system and many [planets](#) outside of it, but to date, no one has captured evidence of a moon orbiting one of those exoplanets. This might change, as the team studying data from the Kepler Space Telescope believe they have found strong evidence for a moon orbiting a planet which is itself orbiting a star called Kepler-1625.

Exoplanets are found by noting the dimming of [stars](#) that occurs as a planet passes between Earth and the star. A moon would be found in roughly the same way, the researchers note, by looking for dimming that occurs in the light reflected from a planet caused by the transit of a moon. Kipping and his team report that they recorded three such dippings as the planet made three trips around its star. They have given their find a statistical confidence of slightly above 4 sigma, noting that the dipping they observed could be chalked up to a fluke in the data—there also exists the possibility that the dimming has another cause. They note that the star system is approximately 4000 light years away, which means the light from the planet is extremely dim to begin with.

The team expects their finding to be verified (or quashed) by data from the Hubble Space Telescope, which offers much better data, sometime in the near future. But while they wait, they are already developing theories about the exomoon—if it truly exists, it would have to be much larger than our moon in order to detect it, perhaps as large as Neptune. That would mean the planet it is orbiting was also quite large, likely as large as Jupiter. If that is the case, its size suggests that it would likely have formed later than moons in our solar system. It would also make the first observed exomoon the largest [moon](#) ever observed.

More information: HEK VI: On the Dearth of Galilean Analogs in Kepler and the Exomoon Candidate Kepler-1625b I, arXiv:1707.08563 [astro-ph.EP] arxiv.org/abs/1707.08563

Abstract

Exomoons represent an outstanding challenge in modern astronomy, with the potential to provide rich insights into planet formation theory and habitability. In this work, we stack the phase-folded transits of 284 viable moon hosting Kepler planetary candidates, in order to search for satellites. These planets range from Earth-to-Jupiter sized and from ~ 0.1 to 1.0 AU in separation - so-called "warm" planets. Our data processing includes two-pass harmonic detrending, transit timing variations, model selection and careful data quality vetting to produce a grand light curve with a r.m.s. of 5.1 ppm. We find that the occurrence rate of Galilean-analog moon systems can be constrained to be η

Citation: Possible first sighting of an exomoon (2017, July 28) retrieved 27 April 2024 from <https://phys.org/news/2017-07-sighting-exomoon.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.