


# A new model yields insights into glaciers' retreats and advances

July 25 2017, by Meghan Murphy



The terminus of the Taku Glacier, in the images at left, advanced more than three miles from 1933 to 2016, as reflected by the yellow and blue lines superimposed on the photographs. The location of the red dot on a mountain in the top image matches the location of the righthand red dot in the lower image. The images and lines at right show the Columbia Glacier's retreat from 1985 to 2016. Credit: Image courtesy of Douglas Brinkerhoff, from photography by the US Navy, the National Snow and Ice Data Center, Martin Truffer and the

Landsat 7 and 8 satellites via US Geological Survey


A University of Alaska Fairbanks study looking at the physics of tidewater glaciers has yielded new insights into what drives their retreat-and-advance cycles and the role that climate plays in these cycles.

Lead author and UAF geophysics doctoral student Douglas Brinkerhoff said the study in *Nature Communications* reveals that shifting sediments drive the cycles among [tidewater glaciers](#) in temperate climates such as southern Alaska.

The study also reveals that these [glaciers](#) don't need periods of warming within the temperate climates to trigger the glacier's retreat, as previously thought.

"Tidewater glaciers can advance further into the ocean by riding on top of a pile of their own [sediment](#), but this can get them into trouble," said Brinkerhoff. "Eventually the glacier snout slows down, but it continues to push that sediment pile further out to sea, essentially pulling the rug out from under itself. When the glacier floats, there's no resistance at its base holding it back and the whole floating part tends to disintegrate."

The findings stem from a mathematical model that Brinkerhoff developed to better understand the periodic retreats and advances of glaciers that flow into the ocean. One-third of Alaska's approximately 60 tidewater glaciers are advancing as part of this cycle, despite the otherwise widespread glacier loss. The advances usually last several centuries, while the retreats take only decades.



This illustration shows a tidewater glacier slowly advancing on a sediment pile. The melting glacier terminus eventually begins to erode the sediment pile. That undercuts the ice's support and triggers a rapid collapse, in a process explained by a new model developed by a University of Alaska Fairbanks researcher. Credit: Illustration by Meghan Murphy

Co-author Martin Truffer, a glaciologist with UAF's Geophysical Institute, said that glaciologist Austin Post first noted the tidewater glacier cycles in the 1970s. He saw that sediment piles developed in the ocean in front of advancing glaciers. These piles allowed the glaciers to extend their advance, followed by often catastrophic collapse and retreat.

"While the work of Austin Post and others clearly showed that the erosion, evacuation and deposition of glacial sediments play an important role, the work reported here managed, for the first time, to capture all the relevant processes in one single model," said Truffer.

Brinkerhoff said that many scientists believed that warming periods within the temperate climate triggered the extended glacier to collapse. They thought that the extended part of the glacier was vulnerable to slight periods of warming because it was flatter and at sea level.

But the model shows that the extended portion of the glacier would collapse even without these warming periods because the glacier erodes the pile supporting it.





Martin Truffer drills holes for explosives in the surface of Taku Glacier. The explosions produce seismic waves that bounce off the sediment below the glacier, providing information about its thickness and character. Credit: Douglas Brinkerhoff

Co-author Andy Aschwanden, a modeler with UAF's Geophysical Institute, said that tidewater glaciers in climates colder than Alaska, such as southeastern Greenland or the northern Antarctic Peninsula, generally do not exhibit a [cycle](#) yet.

"The model suggests if temperatures keep rising some of the tidewater glaciers in colder areas may start to advance, even though that seems counterintuitive," Aschwanden said.

To develop and test the model, Brinkerhoff used observations from glaciers such as Prince William Sound's Columbia Glacier, which started retreating in 1985. He said they also looked at advancing tidewater glaciers such as Hubbard Glacier, which has a submarine pile in front of it, and Taku Glacier near Juneau, the terminus of which now rests on an exposed pile of sediment.

**More information:** Douglas Brinkerhoff et al, Sediment transport drives tidewater glacier periodicity, *Nature Communications* (2017). [DOI: 10.1038/s41467-017-00095-5](https://doi.org/10.1038/s41467-017-00095-5)

Provided by University of Alaska Fairbanks

Citation: A new model yields insights into glaciers' retreats and advances (2017, July 25)  
retrieved 17 May 2024 from <https://phys.org/news/2017-07-yields-insights-glaciers-retreats-advances.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.