


# New model for bimolecular reactions in nanoreactors

August 4 2017

---



Sketch of an yolk-shell-type nanoreactor: reactands A and B diffuse through the shell and react to C at the catalytically active nanoparticle (yellow). Credit: HZB

Theoretical physicists have devised a mathematical model of two different molecules reacting within so called nanoreactors that act as catalysts. They gained surprising new insights as to what factors promote reactions and how to control and select them. The model is relevant for a wide range of research fields, from biophysics to energy materials.

Nanoreactors are tiny systems which facilitate specific [chemical reactions](#), as a catalyst does. Many are found in biological systems, such as certain proteins. But chemists are also able to synthesise artificial nanoreactors to control chemical reactions. An important class of these nanoreactors has a "yolk & [shell](#)" architecture like an egg: a catalytically active metallic nanoparticle is surrounded by a shell consisting of a polymeric network. These kinds of nanoreactors can create isolated environments for specific reactions and restrict them to the tiny space inside the shell.


## **Mathematical description delivers new insights**

"We have now mathematically described for the first time how two different molecules are transported to react within nanoreactors. The [new model](#) shows clearly what factors favour a given [reaction](#)", says Dr. Rafael Roa. Roa is first author of the publication in *ACS Catalysis* and a postdoc in the group headed by Prof. Joe Dzubiella at the HZB Institute for Soft Matter and Functional Materials.

## **What matters most?**

Some of the results come as a surprise: contrary to expectations, the reaction rate is not so much limited by the concentration of the molecules in solution, but decisively by the permeability of the nanoreactor's shell. "This is extremely interesting since chemists today can often fine tune or even switch the permeability of these shells to

specific molecules via variations in temperature or other parameters", explains co-author Dr. Won Kyu Kim.



Reactants A and B diffuse through the shell and react to product C at the catalytically active nanoparticle (yellow) inside. Credit: HZB

### Photo-activation taken into account

The new [model](#) is a big step forward from the older theory done many decades earlier that could handle only one molecule. "Our model is applicable to research on [energy materials](#), and it can even take into account photo-activation of one of the [molecules](#) at the shell by

sunlight", Dzubiella states. He has achieved with this work one of the goals of his European Research Council (ERC) Consolidator Grant (2015-2020).

## Predictions will be put to test

Dzubiella's Soft Matter Theory group collaborates with HZB chemist Prof. Yan Lu, an acknowledged expert in synthetic nanoreactors. They are eager to test their theoretical predictions on real systems. "We're now able to better understand what happens, and we expect to predict how the catalytic effects of these types of nanoreactors can be controlled - through feedback loops, for instance, which will stop or start the reaction at will."

**More information:** Rafael Roa et al, Catalyzed Bimolecular Reactions in Responsive Nanoreactors, *ACS Catalysis* (2017). [DOI: 10.1021/acscatal.7b01701](https://doi.org/10.1021/acscatal.7b01701)

Provided by Helmholtz Association of German Research Centres

Citation: New model for bimolecular reactions in nanoreactors (2017, August 4) retrieved 19 April 2024 from <https://phys.org/news/2017-08-bimolecular-reactions-nanoreactors.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--