

Biodiversity negates carbon storage in seagrasses, new study finds

August 7 2017, by Evelyn S. Gonzalez



FIU researchers take samples of deep sediments in Florida Bay. Credit: Florida International University

Seagrass meadows are great absorbers of carbon dioxide from the air. But the algae, animals, corals and plants that live among them release large amounts of carbon dioxide, according to newly released research.



The scientists are now identifying seagrass locations with fewer emitters to target for conservation.

Scientists at Florida International University examined <u>seagrass</u> <u>meadows</u> in Florida Bay, some of the largest on Earth, where waters are warm and plant and animal abundance is high. They compared these ecosystems to those in southeastern Brazil where meadows are smaller, waters are cooler, and plant and animal abundance is lower. They found, although Florida Bay's <u>seagrasses</u> act as carbon sinks, the organisms living among them offset the benefits of seagrass carbon storage by releasing <u>carbon dioxide</u>.

"In seagrass meadows, these two processes happen simultaneously and have opposite effects on carbon sequestration," said Jason Howard, researcher in FIU's Marine Education Research Initiative and lead author of the study. "If we want to mitigate the most <u>carbon dioxide emissions</u>, we need to understand these competing processes and choose conservation sites accordingly."

The world's seagrasses are disappearing at alarming rates because of grazing, overfishing, pollution and poor management. The findings could aid decision-makers in identifying seagrass meadows with greatest overall carbon dioxide storage potential, so they can target conservation efforts and maximize sequestration.

Scientists studying seagrasses have traditionally focused on the role of organic carbon in the carbon storage cycle. As seagrasses turn sunlight into energy, they take in carbon dioxide to make plant tissue. When they die, the tissue decomposes and organic carbon accumulates in the soil. This organic material is the basis of carbon storage inventories and mitigation plans. At the same time, the organisms living in seagrass beds build their skeletons and shells with calcium carbonate, which is made up of inorganic carbon. When the inorganic carbon is considered in



carbon storage inventories and mitigation plans, the seagrass ecosystems of Brazil store more carbon dioxide than those in Florida Bay.

"The results of this study were totally unexpected," said James Fourqurean, director of FIU's Marine Education Research Initiative and co-author of the study. "They could influence the way seagrass carbon stores are inventoried at the national level, as required by international climate change mitigation treaties."

Fourqurean has dedicated his career to seagrasses. His research provided the first global analysis of carbon stored in seagrasses. He has presented worldwide and testified before the European Union championing for Blue Carbon, a global initiative that allows regulated sources to buy credits for greenhouse emissions, and help restore and preserve seagrasses for mitigation. The money is used to preserve the world's seagrasses, mangroves, marshes and other resources that naturally store carbon.

Most seagrass research is done in areas with big seagrass meadows, including south Florida, the Mediterranean and southeast Asia. South America has been overlooked because of its lack of seagrass coverage and species diversity. Since the ability of seagrasses to store carbon varies from place to place, the researchers call for more thorough studies on the relationship between seagrasses, the local environment and the carbon cycle.

FIU's Marine Education Research Initiative is housed in the Institute of Water and Environment. FIU launched the institute in 2016 to address global water and environmental issues. It brings together some of FIU's top centers and programs to expand research and community engagement in the face of growing environmental threats, including the Center for Aquatic Chemistry and Ecotoxicology, Sea Level Solutions Center and Southeast Environmental Research Center, as well as



Everglades programs and international water programs.

The study was recently published in *Limnology and Oceanography*. It was done in collaboration with the State University of Rio de Janeiro in Brazil.

More information: Jason L. Howard et al. CO2 released by carbonate sediment production in some coastal areas may offset the benefits of seagrass "Blue Carbon" storage, *Limnology and Oceanography* (2017). DOI: 10.1002/lno.10621

Provided by Florida International University

Citation: Biodiversity negates carbon storage in seagrasses, new study finds (2017, August 7) retrieved 27 June 2024 from <u>https://phys.org/news/2017-08-biodiversity-negates-carbon-storage-seagrasses.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.