

# Innovative carbon nanotube photocatalytic materials for efficient solar energy conversion and hydrogen production

August 24 2017




Figure 1: Absorption spectrum of s-SWCNTs (black line) overlaid with solar spectrum (yellow line). Credit: Okayama University

The unique properties of semiconducting single-walled carbon nanotubes

(s-SWCNTs) offer significant advantages over organic molecules, semiconducting polymers, and solid-state semiconductors for wide ranging applications. In particular, s-SWCNT are potentially highly effective active absorption layers in thin-film solar cells because the optical absorption bands—that depend on the chiral indices,  $(n,m)$ —of s-SWCNTs exhibit excellent overlap with the useful solar spectrum range of the sun's radiation.

However, in other applications of carbon nanotubes, there have not been any reports on the use of s-SWCNTs as components of photocatalysts for [splitting water](#) into hydrogen (photocatalytic H<sub>2</sub>) although photocatalytic-based water splitting is expected to be a key technology for [solar energy conversion](#) and sustainable production of hydrogen.

Now, Yutaka Takaguchi and colleagues at Okayama University, Yamaguchi University, and Tokyo University of Science report on the observation of photocatalytic H<sub>2</sub> evolution from water triggered by photoexcitation of s-SWCNTs.

The researchers fabricated a structure consisting of a s-SWCNT/C60 coaxial heterojunction by a self-organization technique using fullerodendron to make s-SWCNT act as a photocatalyst. This heterojunction was used to induce the highly efficient H<sub>2</sub> evolution reaction from water, where the (8, 3) SWCNT/fullerodendron coaxial photocatalyst shows H<sub>2</sub>-evolving activity ( $QY = 0.015$ ) upon 680-nm illumination, which is E22 absorption of (8, 3) SWCNT.

Due to strong absorption coefficients and ease of modifying s-SWCNTs, the CNT-photocatalyst could be a powerful candidate as a material for solar energy conversion and H<sub>2</sub> production without CO<sub>2</sub> emission.




Figure 2: Illustration of SWCNT photocatalysts consisting of the s-SWCNT/C<sub>60</sub> coaxial heterojunction. Credit: Okayama University

**More information:** Noritake Murakami et al. SWCNT Photocatalyst for Hydrogen Production from Water upon Photoexcitation of (8, 3) SWCNT at 680-nm Light, *Scientific Reports* (2017). [DOI: 10.1038/srep43445](https://doi.org/10.1038/srep43445)

Provided by Okayama University

Citation: Innovative carbon nanotube photocatalytic materials for efficient solar energy conversion and hydrogen production (2017, August 24) retrieved 19 April 2024 from <https://phys.org/news/2017-08-carbon-nanotube-photocatalytic-materials-efficient.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.