

Cockroach gardeners: Spreading plant seeds across the forest floor

August 4 2017



This heterotrophic herb lacks chlorophyll making its appearance almost all white. Credit: Yasuhiro Uehara and Naoto Sugiura

In forest ecosystems, cockroaches are known as important decomposers that consume dead and decaying plants. Quite unexpectedly, however, researchers have found that they also provide seed dispersal services for the plant *Monotropastrum humile*, a forest-floor herb belonging to the azalea family (Ericaceae). This entirely new mode of plant-insect interaction is reported online in the July 27th, 2017 issue of the *Botanical Journal of the Linnean Society*.

M. humile is mycoheterotrophic, which means that it completely relies on Russulaceae fungi for nutrients. Its fleshy [fruit](#) contains hundreds of seeds that are very small (ca. 0.3 mm long and 0.2 mm wide) and have a hard testa (seed coat). As the fruit ripens, they drop to the ground or the whole plant stalk falls down, making the pulp available at ground level. The ripe fruit has no distinctive scent, are dull white in color, and are inconspicuous against the litter layer of the shady [forest floor](#). The slightly juicy pulp is not sweet to the human palate.

The fruit attracted various invertebrates but birds and mammals seemed uninterested. Of the invertebrates, only the [forest](#) cockroach *Blattella nipponica* (mostly adults) consistently visited and consumed the seed-embedded pulp at night. Adult *B. nipponica* excreted feces about three to 10 hours after their collection from the field, and each fecal pellet (ca. 1 mm long) contained an average of 3.1 *M. humile* seeds.

Seeds in the feces remained intact and viable. A chemical test revealed that about half of the cockroach-ingested seeds remained viable and that their viability (52.0 percent) was not significantly different from that of seeds directly removed from the pulp (49.3 percent). This indicates that *B. nipponica* cockroaches can function as a legitimate seed disperser for *M. humile*.



Japanese endemic cockroaches (*B. nipponica*) feeding on the fruit of the *M. humile* plant. It will ingest the seeds of the plant and disperse them later across the forest floor. Credit: Yasuhiro Uehara and Naoto Sugiura

M. humile [plants](#) have traits that may reflect cockroach-mediated seed dispersal. First, their fruit ripening period largely coincides with the eclosion period (May and June) of the forest cockroach. Second, the presentation of its fruit at ground level where the forest cockroaches inhabits. Third, the small seeds with a hard testa enables ingested seeds to remain viable. Furthermore, *B. nipponica* adults have intrinsic traits that may facilitate their actions as dispersal agents for this mycoheterotrophic herb: (1) their abundance in the plant habitat; (2)

they are ground-dwelling, which means they defecate on the forest floor where *M. humile*'s fungal partners, Russulaceae, reside; and (3) the long transit times of seeds through their gut (three to 10 h) providing occasional long-distance seed dispersal. Consequently, the findings strongly suggest that the fleshy fruit-bearing *M. humile* is considerably specialized for seed dispersal by adult cockroaches, enabling the plant to steadily disperse minute seeds across the forest floor.

"Given that about 4600 cockroach species have been identified to date, cockroach-mediated seed dispersal may be a more pervasive but as yet undocumented mechanism, rather than a unique exploit by a particular plant species," said Associate Professor Naoto Sugiura of Kumamoto University. "It is hoped that the findings reported here will encourage people to pay more attention to plant-cockroach interactions and seek out further examples of [cockroach](#)-mediated [seed dispersal](#)."

This finding was posted online in the *Botanical Journal of the Linnean Society* on 27 July 2017.



A cockroach (*B. nipponica*) fecal pellet, approximately 1 mm in length, containing intact *M. humile* seeds. The plant is distributed across the forest floor with the help of these insects. Credit: Yasuhiro Uehara and Naoto Sugiura

More information: Yasuhiro Uehara et al, Cockroach-mediated seed dispersal in *Monotropastrum humile* (Ericaceae): a new mutualistic mechanism, *Botanical Journal of the Linnean Society* (2017). [DOI: 10.1093/botlinnean/box043](https://doi.org/10.1093/botlinnean/box043)

Provided by Kumamoto University

Citation: Cockroach gardeners: Spreading plant seeds across the forest floor (2017, August 4)
retrieved 26 April 2024 from

<https://phys.org/news/2017-08-cockroach-gardeners-seeds-forest-floor.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.