

Running out of gas: Gas loss puts breaks on stellar baby boom

August 2 2017





Gas rich galaxies detected with ALMA are shown in red and marked with circles. Most gas rich galaxies are located in the outer part, not the center, of the galaxy cluster (around the center of the image). Credit: ALMA (ESO/NAOJ/NRAO), Hayashi et al., the NASA/ESA Hubble Space Telescope

Understanding the history of star formation in the Universe is a central theme in modern astronomy. Various observations have shown that the star formation activity has varied through the 13.8 billion-year history of the Universe. The stellar birthrate peaked around 10 billion years ago, and has declined steadily since then. However, the cause of the declining stellar birthrate is still not well understood.

"Aiming to investigate what suppresses the <u>star formation</u> activity, we focused on the environment around the galaxies," said Masao Hayashi at the National Astronomical Observatory of Japan (NAOJ).

Hayashi and his colleagues observed the galaxy <u>cluster</u> XMMXCS J2215.9-1738 located 9.4 billion light-years away with the Atacama Large Millimeter/submillimeter Array (ALMA). Because it takes time for the light from distant objects to reach us, observing far-away galaxies shows us what the Universe looked like when the light was emitted. In this case, the light from XMMXCS J2215.9-1738 was emitted 9.4 billion years ago, which is around the time that the stellar birthrate peaked. In fact, previous observations with NAOJ's Subaru Telescope revealed that many of the galaxies in the cluster are actively forming <u>stars</u>.

ALMA detected radio signals emitted from <u>carbon monoxide gas</u> in 17 of the galaxies in the cluster. This is a record-high number for the detection of gas-rich galaxies at such a distance. Interestingly, the gasrich galaxies detected with ALMA are located towards the outer part of



the galaxy cluster, not in the center. This is the first time ever that such a location differentiation has been found in a galaxy cluster 10 billion light-years away.

The team assumes that the gas-rich galaxies detected with ALMA are in an intermediate step in the process of becoming members of the cluster. As new member galaxies pass through the hot gas filling the cluster, cold gas in the galaxies is stripped away by the hot gas. Active star formation consumes what little gas survives in the galaxies. As the cold gas needed to make stars runs out, star formation stops.

Actually, there are some galaxies with active star formation at the central part of the cluster. The team suggests that they are rather evolved, old members of the cluster consuming the last of their gas to form stars.

"Recent observational and theoretical studies show that the distribution of gas is key to understanding the evolution of galaxies," explains Hayashi. "Our observations provide robust statistics showing that a number of gas-rich galaxies are located in the outer part of a <u>galaxy</u> <u>cluster</u>. With this result we have opened a future path for revealing the evolutionary process of <u>galaxies</u> in galaxy clusters."

These observation results were published as Hayashi et al. "Evolutionary Phases of Gas-rich Galaxies in a Galaxy Cluster at z = 1.46" in the *Astrophysical Journal Letters* in May 2017.

More information: Masao Hayashi et al, Evolutionary Phases of Gasrich Galaxies in a Galaxy Cluster at z = 1.46, *The Astrophysical Journal* (2017). DOI: 10.3847/2041-8213/aa71ad

Provided by National Institutes of Natural Sciences



Citation: Running out of gas: Gas loss puts breaks on stellar baby boom (2017, August 2) retrieved 2 May 2024 from <u>https://phys.org/news/2017-08-gas-loss-stellar-baby-boom.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.