


Physicists explore a new recipe for heating plasma

August 22 2017, by Paul Rivenberg



The interior of the Alcator C-Mod tokamak, where experiments were conducted that have helped create a new scenario for heating plasma and achieving fusion. Credit: Bob Mumgaard/Plasma Science and Fusion Center

In the quest for fusion energy, scientists have spent decades

experimenting with ways to make plasma fuel hot and dense enough to generate significant fusion power. At MIT, researchers have focused their attention on using radio-frequency (RF) heating in magnetic confinement fusion experiments like the Alcator C-Mod tokamak, which completed its final run in September 2016.

Now, using data from C-Mod experiments, [fusion](#) researchers at MIT's Plasma Science and Fusion Center (PSFC), along with colleagues in Belgium and the UK, have created a new method of heating fusion plasmas in tokamaks. The new method has resulted in raising trace amounts of ions to megaelectronvolt (MeV) energies—an order of magnitude greater than previously achieved.

"These higher [energy](#) ranges are in the same range as activated fusion products," PSFC research scientist John C. Wright explains. "To be able to create such energetic ions in a non-activated device—not doing a huge amount of fusion—is beneficial, because we can study how ions with energies comparable to fusion reaction products behave, how well they would be confined."

The new approach, recently detailed in the journal *Nature Physics*, uses a fuel composed of three ion species: hydrogen, deuterium, and trace amounts (less than 1 percent) of helium-3. Typically, [plasma](#) used for fusion research in the laboratory would be composed of two ion species, deuterium and hydrogen or deuterium and He-3, with deuterium dominating the mixture by up to 95 percent. Researchers focus energy on the minority species, which heats up to much higher energies owing to its smaller fraction of the total density. In the new three-species scheme, all the RF energy is absorbed by just a trace amount of He-3 and the ion energy is boosted even more—to the range of activated fusion products.

Wright was inspired to pursue this research after attending a lecture in

2015 on this scenario by Yevgen Kazakov, a researcher at the Laboratory for Plasma Physics in Brussels, Belgium, and the lead author of the Nature Physics article. Wright suggested that MIT test these ideas using Alcator C-Mod, with Kazakov and his colleague Jef Ongena collaborating from Brussels.

At MIT, PSFC research scientist Stephen Wukitch helped developed the scenario and run the experiment, while Professor Miklos Porkolab contributed his expertise on RF heating. Research scientist Yijun Lin was able to measure the complex wave structure in the plasma with the PSFC's unique phase contrast imaging (PCI) diagnostic, which was developed over the last two decades by Porkolab and his graduate students. Research scientist Ted Golfinopoulos supported the experiment by tracking the effect of MeV-range ions on measurements of plasma fluctuations.

The successful results on C-Mod provided proof of principle, enough to get scientists at the UK's Joint European Torus (JET), Europe's largest fusion device, interested in reproducing the results. Like JET, C-Mod operated at magnetic field strength and plasma pressure comparable to what would be needed in a future fusion-capable device. The two tokamaks also had complementary diagnostic capabilities, making it possible for C-Mod to measure the waves involved in the complex wave-particle interaction, while JET was able to directly measure the MeV-range particles.

John Wright praises the collaboration.

"The JET folks had really good energetic particle diagnostics, so they could directly measure these high energy ions and verify that they were indeed there," he says. "The fact that we had a basic theory realized on two different devices on two continents came together to produce a strong paper."

Porkolab suggests that the new approach could be helpful for MIT's collaboration with the Wendelstein 7-X stellarator at the Max Planck Institute for Plasma Physics in Greifswald, Germany, where research is ongoing on one of the fundamental physics questions: How well fusion-relevant [energetic ions](#) are confined. The Nature Physics article also notes that the experiments could provide insight into the abundant flux of He-3 ions observed in solar flares.

More information: Ye. O. Kazakov et al. Efficient generation of energetic ions in multi-ion plasmas by radio-frequency heating, *Nature Physics* (2017). [DOI: 10.1038/nphys4167](https://doi.org/10.1038/nphys4167)

Provided by Massachusetts Institute of Technology

Citation: Physicists explore a new recipe for heating plasma (2017, August 22) retrieved 23 April 2024 from <https://phys.org/news/2017-08-physicists-explore-recipe-plasma.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--