

# Can poor air quality mask global warming's effects?

August 8 2017

---



Looking through smog in downtown Atlanta from midtown. Credit: CC BY-SA 2.0, by Flickr user Ben Ramsey

During the 20th century, the average temperature of the continental United States rose by almost 1 degree Fahrenheit (0.5 degree Celsius)—everywhere, that is, except in the Southeast. There, until the 1980s, the temperature actually decreased slightly. Climate scientists dubbed this peculiar phenomenon the "warming hole," and it was the cause of much speculation. But beginning in the 1990s, temperatures in the Southeast began to warm again, and in the early years of the 21st century this warming has accelerated.


A new study published in the journal *Remote Sensing* presents evidence that a significant improvement in air quality in the region may have contributed to the disappearance of the warming hole after about 1990—and that other polluted regions outside the United States, such as China and India, may experience the same phenomenon.

One major factor in [poor air quality](#) is airborne aerosols—tiny particles of dust, soot from wood burning, coal and oil combustion, or sulfates created by precursor gases emitted from factories and car exhaust, to name a few sources. Aerosols can decrease temperature by dimming sunlight at Earth's surface and by increasing the amount and lifetimes of clouds, which reflect sunlight back into space.

After the warming hole mysteriously disappeared, various studies proposed possible causes: changes in cloud cover, precipitation or in the amount of aerosols produced by air pollution. In 2006, the U.S. Environmental Protection Agency (EPA) began implementing a more stringent cap on the concentration of aerosol particles smaller than about 1/10,000th of an inch (2.5 micrometers) in diameter. To comply with the regulation, many U.S. power utilities and industrial companies began reducing their use of coal and installing filters to reduce emissions.

A similar change to temperature trends occurred in Europe in the 1980s after new regulations improved air quality there. Because reduced

aerosol particle concentrations allow more sunlight to reach Earth's surface, the scientists hypothesized that the improvements in U.S. [air quality](#) could also be responsible for the [temperature change](#) over the Southeast.



U.S. trends in the optical depth of aerosols from 2001 to 2015 (blue is decreasing, red is increasing). Improvements in regional air quality in the Southeast U.S. may have contributed to warmer temperatures seen in the region since about 1990. Credit: NASA-JPL/Caltech/School of the Art Institute of Chicago

To test this hypothesis, a team led by Mika Tosca, a researcher at NASA's Jet Propulsion Laboratory in Pasadena, California (who is now with the School of the Art Institute of Chicago), used three surface

temperature data sets. The data sets were compiled by the University of Delaware, the University of California (UC) at Berkeley, and the Global Historical Climatology Network (which compiles surface temperature and precipitation data). They also used [aerosol](#) data from two satellite instruments: the Multi-angle Imaging SpectroRadiometer (MISR) instrument on NASA's Terra satellite, launched in 1999, and the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) on the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) satellite, a joint mission between NASA and the French space agency, CNES, launched in 2006.

The data show that between 2000 and 2015, while summertime temperatures in the Southeast United States increased by roughly 1.5 degrees Fahrenheit (0.75 degree Celsius), significantly faster than the increase in the continental United States during the 20th century, the amount of summertime aerosols decreased overall by about 20 percent, with a much steeper decline after 2007. The timing of this decline coincided with the implementation of the new EPA standards.

To help determine how much of the temperature change was caused by the changes in aerosols, Tosca and colleagues used a model that simulates how the sun's energy travels through Earth's atmosphere, using the MISR and CALIOP satellite data as inputs. The increase in sunlight shown in the model results matches well with daily measurements taken at a National Oceanic and Atmospheric Administration (NOAA) solar radiation monitoring station in Goodwin Creek, Mississippi, suggesting that the decrease in aerosols is a plausible explanation for most of the disappearance of the warming hole.

Tosca acknowledges that linkages between aerosols and clouds could also play a role. The next step would be to run a more sophisticated climate model that takes into account clouds and the aerosols' effects on them. The team would also like to apply this kind of analysis to other

areas with high air pollution levels, such as China and India. They hypothesize that these areas might have "warming holes" of their own—regions where the effects of climate change are being muted by the high concentrations of aerosols in the atmosphere. If these areas reduce [air pollution](#) in the future, they might experience a sudden [temperature](#) jump as well.

"Overall, the goal is to more accurately predict what will happen to our planet," Tosca said. "This type of observation-based research gives us better models, better models give us better forecasts, and better forecasts enable better policy."

The study is titled "Attributing Accelerated Summertime Warming in the Southeast United States to Recent Reductions in Aerosol Burden: Indications from Vertically-Resolved Observations."

**More information:** Mika Tosca et al. Attributing Accelerated Summertime Warming in the Southeast United States to Recent Reductions in Aerosol Burden: Indications from Vertically-Resolved Observations, *Remote Sensing* (2017). [DOI: 10.3390/rs9070674](https://doi.org/10.3390/rs9070674)

Provided by NASA

Citation: Can poor air quality mask global warming's effects? (2017, August 8) retrieved 8 May 2024 from <https://phys.org/news/2017-08-poor-air-quality-mask-global.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--