


Rare metals in the Himalayas

August 30 2017



Simplified geological map showing the distribution of Himalayan leucogranites. Credit: Science China Press


Two sub-parallel belts of Cenozoic aged Himalayan leucogranite on the Tibetan Plateau extend east to west over more than 1000 km, and are regarded as the largest granitic belts in the world. Rare-metal mineralization was identified in relation to these leucogranites.

The Himalayan leucogranite is unique, with petrological characteristics similar to well-known rare-metal granites worldwide. However,

relatively few studies on the rare-metal mineralization in this region have been published. The research groups in Nanjing University and Institute of Geology and Geophysics, Chinese Academy of Science, organized a field expedition in south Tibet in the summer of 2016 to constrain the distribution of mineralization in the region, which was published in the *Science China Earth Sciences*.

The first discovery identified in the field is the widespread Be-mineralization containing in most of leucogranitic plutons (Fig. 2a). The authors write, "Aquamarine, a variety of beryl, major precious mineral resource in Nepal, has been widely explored, exploited and traded for a long time, and they may also be Be-mineralization potential on the Chinese side of the Himalayas."

Detailed microscope observation and microprobe analysis were conducted at the laboratories and 12 leucogranite plutons were found to contain rare-metal bearing minerals such as beryl (the representative of Be mineralization), columbite-group minerals (Fig. 2b), tapiolite, pyrochlore-microlite, fergusonite, Nb-Ta rutile (the representative of Nb-Ta mineralization), and cassiterite (the representative of Sn mineralization).



(a) These are beryl crystals observed in the outcrops. Euhedral, columnar beryl crystals are marked by the red circles. (b) BSE image of columbite with oscillatory zoning from the Himalaya. Credit: Science China Press

Based on the analytical results, the researchers revealed the distribution of the mineralization as: "Rare-metal mineralization was observed in both the Tethyan and Higher Himalayan belts. No clear differences between the two belts were identified. However differences are clear when comparing the eastern and western parts of the belts. The eastern plutons are characterized by Nb, Ta, Sn, and Be mineralization and Sn mineralization is notably absent in western plutons. These results suggest that rare-metal mineralization in the Himalayan leucogranites is regionally variable, but does not appear to be controlled by tectonic characteristics of granite emplacement and the pluton size."

Petrogenesis was also prepared for the Himalayan rare-metal leucogranite. The model of "magmatic fractionation" and the abundance of fluxing components in the melt (e.g., H₂O, Li, F, B, and P) are very important for the formation of the granite and enrichment of rare metal in the melt and/or fluids.

Rare metal has the widespread application value in the strategic development of new industries. "China ranks high in resources and production of rare-metal mineral, especially in two granitic belts in the Nanling range and the Altai district (Xinjiang) with several world-class deposits", the authors in the article pointed out the immense value of the investigation in the Himalaya: "Our preliminary study on the Himalayan region shows additional enormous potential for the rare-metal mineralization in what may become the country's economically important metallogenic belt."

More information: RuCheng Wang et al, A preliminary study of rare-metal mineralization in the Himalayan leucogranite belts, South Tibet, *Science China Earth Sciences* (2017). [DOI: 10.1007/s11430-017-9075-8](https://doi.org/10.1007/s11430-017-9075-8)

Provided by Science China Press

Citation: Rare metals in the Himalayas (2017, August 30) retrieved 29 January 2023 from <https://phys.org/news/2017-08-rare-metals-himalayas.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.