

New dust sources from a shrinking Salton Sea have negative ecological and health impacts

August 3 2017







The Salton Sea from space. Credit: NASA

Scientists at the University of California, Riverside investigating the composition of particulate matter (PM) and its sources at the Salton Sea have found that this shrinking lake in Southern California is exposing large areas of dry lakebed, called playa, that are acting as new dust sources with the potential to impact human health.

"Playas have a high potential to act as dust sources because playa surfaces often lack vegetation," said Roya Bahreini, an associate professor of environmental sciences, who led the research project. "Dust emissions from playas increase airborne PM mass, which has been linked to cardiovascular disease, respiratory disease, and mortality."

Study results appeared recently in *Environmental Science and Technology*.

Bahreini's team set out to test whether emissions from playas change the composition of PM10 (particulate matter with diameters up to 10 microns) near the Salton Sea. The team assessed the composition of playa soils (recently submerged underneath the Salton Sea), desert soils (located farther from the sea), and PM10 collected during August 2015 and February 2016.

They found that dust sources contributed to about 45 percent of PM10 at the Salton Sea during the sampling period while playa emissions contributed to about 10 percent. Further, they found that playa emissions significantly increased the <u>sodium content</u> of PM10.



"Increase in the sodium content of PM10 can affect the ecosystem when the sodium-rich particles deposit downwind and change the natural balance for soils and agricultural lands or when these particles form clouds," Bahreini said. "Currently, the primary concern for PM emissions from playa is the contribution to total concentration of PM10, which regardless of composition, is an irritant and can have negative respiratory effects."

Her team also found that playa soils and PM10 are significantly enriched in selenium relative to desert soils.

Bahreini explained that selenium can be a driver of aquatic and avian toxicity. "Additionally, higher selenium enrichments in PM10 during summertime suggest that selenium volatilization from the playa may become an important factor controlling the selenium budget in the area as more playa gets exposed," she said.



Alexander L. Frie, a graduate student in environmental sciences and the first



author of the research paper, at the sampling site. Credit: Bahreini lab, UC Riverside.

Alexander L. Frie, a graduate student in environmental sciences and the first author of the research paper, urges that the Salton Sea be paid close attention since, although it is widely considered a large ecological disaster, with no serious monitoring and remediation efforts the sea may also create a <u>human health</u> crisis for the surrounding area.

"Although our results indicate playa emission may not be especially toxic when compared to other dust particles, increased emissions from a shrinking sea will reduce the quality of life of the residents within the Imperial Valley and other downwind regions," he said.

Samantha C. Ying, an assistant professor of environmental sciences and a coauthor on the paper, stresses that monitoring the increase in dust sources over time is necessary to quantify its contribution to local health problems.

"Our study shows that the shrinking Salton Sea is contributing to dust sources in the region," she said. "Even considering just the small area of playa that is exposed now, the contributions are significant."

Another concern the researchers point out is that water that is currently diverted from the Colorado River and directed into the Salton Sea is scheduled to end before 2018. The resultant decrease of inflow into the sea will likely cause a decline in water level, exposing more playa, and therefore emitting more <u>dust</u>.

"With more playa being exposed, we expect total PM10 concentrations to increase and human exposure to these particles in downwind areas will



also increase," Bahreini said. "Therefore implementing any project, for example, creating shallow water pools over the playa, that limits formation of salt crusts on the playa will be valuable."

Bahreini, Frie and Ying were joined in the study by Justin H. Dingle, a graduate student in Bahreini's lab.

More information: Alexander L. Frie et al. The Effect of a Receding Saline Lake (The Salton Sea) on Airborne Particulate Matter Composition, *Environmental Science & Technology* (2017). DOI: <u>10.1021/acs.est.7b01773</u>

Provided by University of California - Riverside

Citation: New dust sources from a shrinking Salton Sea have negative ecological and health impacts (2017, August 3) retrieved 27 April 2024 from <u>https://phys.org/news/2017-08-sources-salton-sea-negative-ecological.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.