

Wild sheep grazed in the Black Desert 14,500 years ago

August 22 2017





Semi-subterranean basalt paved structure in the Black Desert, dated to nearly 14,500 years ago. Credit: University of Copenhagen



Excavations of architecture and associated deposits left by huntergatherers in the Black Desert in eastern Jordan have revealed bones from wild sheep - a species previously not identified in this area in the Late Pleistocene. According to the team of University of Copenhagen archaeologists, who led the excavations, the discovery is further evidence that the region often seen as a 'marginal zone' was capable of supporting a variety of resources, including a population of wild sheep, 14,500 years ago.

A study by a team of archaeologists based at the University of Copenhagen published today in the Royal Society journal *Open Science* documents that the region now known as the Black Desert in eastern Jordan could sustain a population of wild sheep.

"On the basis of morphological and metrical analysis of the faunal remains from Natufian and Pre-Pottery Neolithic A hunter-gatherer deposits, we can document that wild sheep would have inhabited the local environment year-round and formed an important resource for the human population to target for food. Most significantly, however, the presence of the substantial number of bones identified as mouflon extends the known range of wild sheep. This means that we cannot rely on broad scale maps showing ancient wild animal distributions as neat lines," said zooarchaeologist and first-author of the study Lisa Yeomans of the University of Copenhagen.

Adaptive hunter-gatherers

The team have been investigating human occupation in the Late Pleistocene of eastern Jordan. The Levant (i.e. modern-day Jordan, Israel, Palestine, Lebanon and Syria) has long been recognised as an important region associated with changes in social complexity and shifts in subsistence economy that pre-empted the shift to agriculture and farming. Hitherto investigations have generally focused on the Natufian



occupation in the Levantine corridor while eastern Jordan was considered a more marginal environment.

Recent investigations, however, have shown this 'marginal environment' of eastern Jordan to be a resource rich environment which offered people the opportunity to hunt a range of species.

"Our findings illustrate how adaptive humans were nearly 14,500 years ago in a period of climatic change: Wild sheep offered the Natufian and later Pre-Pottery Neolithic populations one of a myriad of resources that could be exploited during the Late Pleistocene even in this more marginal environment beyond the Mediterranean zone. Despite the influences of climate on the resources presented to these hunterforagers, their subsistence strategies were flexible and they could shift focus. Hunting wild sheep is just one of the ways that this is reflected in the archaeological record," said Lisa Yeomans.

More information: Expansion of the known distribution of Asiatic mouflon (Ovis orientalis) in the Late Pleistocene of the Southern Levant, *Royal Society Open Science* (2017). DOI: 10.1098/rsos.170409

Provided by University of Copenhagen

Citation: Wild sheep grazed in the Black Desert 14,500 years ago (2017, August 22) retrieved 2 May 2024 from https://phys.org/news/2017-08-wild-sheep-grazed-black-years.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.