


Dams and other barriers to salmon spawning grounds create challenges for fisheries managers

September 22 2017, by Kat Kerlin



Chinook salmon. Credit: Mike Wier/California Trout

Wild salmon, historically, are born in rivers, swim to sea to live out their adulthoods, and find their way back to their freshwater spawning

grounds to reproduce before dying.

But dams and other barriers to spawning grounds have disconnected that natural cycle, requiring fisheries managers to get creative to support salmon populations. It is now fairly common to transport salmon to and from spawning grounds by truck, boat, and even helicopter.

A report by the University of California, Davis, Center for Watershed Sciences and nonprofit California Trout assesses one such method of assisted movement over dams—two-way trap and haul—being proposed by federal agencies as a high-priority recovery strategy for chinook salmon populations in California.

The report's findings indicate that such programs should proceed with extreme caution, though they may be appropriate in cases where few other options remain to recover dwindling populations of salmon.

What is two-way trap and haul?

Two-way trap and haul involves capturing adult fish below large dams and transporting them to a release site above the dam to spawn. The resulting juvenile fish are then captured, transported and released below the dam, where they will continue to grow or migrate downstream to the ocean.

Fisheries managers in California hope this process helps Central Valley spring-run chinook salmon and Sacramento River winter-run chinook, which are listed, respectively, as threatened and endangered under the federal Endangered Species Act. Most major rivers are dammed in California, which has eliminated access of [wild salmon](#) to most of their historical spawning habitat. This is a major cause of their decline.

"Very few Sacramento River winter-run chinook returned this year,"

said lead author Rob Lusardi, a research ecologist with the UC Davis Center for Watershed Sciences and California Trout. "It was dire. With one more year of drought, they could have blinked out. If two-way trap and haul can successfully reconnect these fish to historical spawning habitat, it would be fantastic. But we need to proceed with caution. The data suggest it's very difficult to get such programs to work effectively to achieve a sustainable population."

Two-way trap and haul has been tested in the Pacific Northwest with uncertain results, and challenges remain.



California's McCloud River is historical spawning habitat for winter-run chinook salmon. The construction of dams eliminated access to that habitat for the

salmon. The fish from these populations were forced to mix and spawn as one population on the Sacramento River. Credit: Peter Moyle/UC Davis

Challenges

The report says the greatest management hurdles include:

- Capturing large numbers of small, out-migrating juvenile salmon before they enter reservoirs.
- The delayed effect of transportation on salmon: While they appear healthy upon release, stress from transport and other factors may increase their risk of death later.
- Long-term sustainability of above-dam populations.
- The unclear influence of hatchery programs on two-way trap-and-haul programs. Salmon numbers may increase when both hatchery and two-way trap-and-haul programs are in place. However, if an increase in adult returns is primarily due to hatchery production of semi-domesticated fish, two-way trap-and-haul programs will contribute little to wild fish production.

How to proceed with caution

If a two-way trap-and-haul project is implemented, the report recommends:

- Clearly defining and measuring success.
- Treating each project as an experiment so the likelihood of success is determined early on.
- Ensuring that the project is part of a broader conservation strategy.

The report concludes that even successful two-way trap-and-haul programs "will not save winter- or spring-run chinook in California but only prolong their decline to extinction" unless they are part of a much more comprehensive program.

"Saving salmon requires managing these amazing fish for their entire life cycle, from eggs in the gravel to adults returning to spawn," said co-author Peter Moyle, professor emeritus with the UC Davis Center for Watershed Sciences.

Such strategies include dam removal, restoring historical floodplain habitat, upgrading water infrastructure, managing the release of water from reservoirs for fish, improving management of fisheries, and improving juvenile survival through the San Francisco Estuary.

High stakes

"We have very carefully examined trap and haul as one of many important strategies in our comprehensive recovery plans for Central Valley salmon," said Maria Rea, assistant regional administrator of NOAA Fisheries' West Coast Region. "The stakes are so great for these highly endangered fish that we cannot afford to pass up this promising option for their recovery. We've lost 90 percent of all Central Valley spawning and rearing habitat behind dams, and trap and haul may be the only way we can help our native fish reclaim those essential rivers and streams."

NOAA Fisheries [recovery plans](#) for endangered Sacramento River winter-run chinook salmon, threatened Central Valley spring-run chinook [salmon](#), and Central Valley steelhead include diverse strategies. These include returning diverted water to spawning streams, restoring historic habitat and improving water quality. They also propose reintroducing the native [fish](#) to spawning streams above dams using a

trap-and-haul approach, and using adaptive management to fine-tune the strategies as biologists learn more.

More information: Robert A. Lusardi et al. Two-Way Trap and Haul as a Conservation Strategy for Anadromous Salmonids, *Fisheries* (2017). [DOI: 10.1080/03632415.2017.1356124](https://doi.org/10.1080/03632415.2017.1356124)

Provided by UC Davis

Citation: Dams and other barriers to salmon spawning grounds create challenges for fisheries managers (2017, September 22) retrieved 3 May 2024 from <https://phys.org/news/2017-09-barriers-salmon-spawning-grounds-fisheries.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--