


Curiosity Mars rover climbing toward ridge top

September 14 2017



Researchers used the Mastcam on NASA's Curiosity Mars rover to gain this detailed view of layers in "Vera Rubin Ridge" from just below the ridge. Credit: NASA/JPL-Caltech/MSSS

NASA's Mars rover Curiosity has begun the steep ascent of an iron-oxide-bearing ridge that's grabbed scientists' attention since before the car-sized rover's 2012 landing.


"We're on the climb now, driving up a route where we can access the layers we've studied from below," said Abigail Fraeman, a Curiosity science-team member at NASA's Jet Propulsion Laboratory in Pasadena, California.

"Vera Rubin Ridge" stands prominently on the northwestern flank of Mount Sharp, resisting erosion better than the less-steep portions of the mountain below and above it. The ridge, also called "Hematite Ridge,"

was informally named earlier this year in honor of pioneering astrophysicist Vera Rubin.

"As we skirted around the base of the ridge this summer, we had the opportunity to observe the large vertical exposure of [rock layers](#) that make up the bottom part of the ridge," said Fraeman, who organized the rover's ridge campaign. "But even though steep cliffs are great for exposing the stratifications, they're not so good for driving up."

The ascent to the top of the ridge from a transition in rock-layer appearance at the bottom of it will gain about 213 feet (65 meters) of elevation—about 20 stories. The climb requires a series of drives totaling a little more than a third of a mile (570 meters). Before starting this ascent in early September, Curiosity had gained a total of about 980 feet (about 300 meters) in elevation in drives totaling 10.76 miles (17.32 kilometers) from its landing site to the base of the ridge.



"Vera Rubin Ridge," a favored destination for NASA's Curiosity Mars rover even before the rover landed in 2012, rises near the rover nearly five years later in this panorama from Curiosity's Mastcam. Credit: NASA/JPL-Caltech/MSSS

Curiosity's telephoto observations of the ridge from just beneath it show finer layering, with extensive bright veins of varying widths cutting through the layers.

"Now we'll have a chance to examine the layers up close as the rover climbs," Fraeman said.

Curiosity Project Scientist Ashwin Vasavada of JPL said, "Using data from orbiters and our own approach imaging, the team has chosen places to pause for more extensive studies on the way up, such as where the rock layers show changes in appearance or composition. But the campaign plan will evolve as we examine the rocks in detail. As always, it's a mix of planning and discovery."




The Mastcam on NASA's Curiosity Mars rover captured this view of "Vera Rubin Ridge" about two weeks before the rover starting to ascend this steep ridge on lower Mount Sharp. Credit: NASA/JPL-Caltech/MSSS

In orbital spectrometer observations, the iron-oxide mineral hematite shows up more strongly at the ridge top than elsewhere on lower Mount Sharp, including locations where Curiosity has already found hematite. Researchers seek to gain better understanding about why the ridge resists erosion, what concentrated its hematite, whether those factors are related, and what the rocks of the ridge can reveal about ancient Martian environmental conditions.

"The team is excited to be exploring Vera Rubin Ridge, as this hematite ridge has been a go-to target for Curiosity ever since Gale Crater was selected as the [landing site](#)," said Michael Meyer, lead scientist of


NASA's Mars Exploration Program at the agency's Washington headquarters.



This view of "Vera Rubin Ridge" from the ChemCam instrument on NASA's Curiosity Mars rover shows sedimentary layers and fracture-filling mineral deposits. Credit: NASA/JPL-Caltech/CNES/CNRS/LANL/IRAP/IAS/LPGN

During the first year after its landing near the base of Mount Sharp, the Curiosity mission accomplished a major goal by determining that billions of years ago, a Martian lake offered conditions that would have been favorable for microbial life. Curiosity has since traversed through a diversity of environments where both water and wind have left their imprint. Vera Rubin Ridge and layers above it that contain clay and sulfate minerals provide tempting opportunities to learn even more about

the history and habitability of ancient Mars.



This view of "Vera Rubin Ridge" from the ChemCam instrument on NASA's Curiosity Mars rover shows sedimentary layers, mineral veins and effects of wind erosion. Credit: NASA/JPL-Caltech/CNES/CNRS/LANL/IRAP/IAS/LPGN

More information: For more about Curiosity, visit mars.jpl.nasa.gov/msl

Provided by Jet Propulsion Laboratory

Citation: Curiosity Mars rover climbing toward ridge top (2017, September 14) retrieved 27 April 2024 from <https://phys.org/news/2017-09-curiosity-mars-rover-climbing-ridge.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.