


# Enzyme's worth to biofuels shown in latest research

September 18 2017

---



NREL scientists Michael Himmel (left), Roman Brunecky, and Yannick Bomble with the X-ray diffraction source used to determine the structure of the *C. bescii* CelA cellulase. Credit: Dennis Schroeder/NREL

An enzyme discovered at the U.S. Department of Energy's (DOE) National Renewable Energy Laboratory (NREL) proves adept at breaking down cellulose fibers regardless of whether their crystalline

structure is simple or highly complex. No other enzyme has shown that ability.

The [enzyme](#), called CelA, comes from *Caldicellulosiruptor bescii*, and NREL scientists reported three years ago, in the journal *Science*, how it can convert biomass to sugars faster than competing catalysts in commercial enzyme preparations. The follow-up study, detailed in the new *Scientific Reports* paper, *The Multi Domain Caldicellulosiruptor bescii CelA Cellulase Excels at the Hydrolysis of Crystalline Cellulose*, points to how the enzyme could help remove one of the technical and economic barriers preventing cellulosic biofuels from becoming a commercial reality.

The [crystalline structure](#) of [cellulose](#) fiber in [plant cell walls](#) generally poses a problem for cellulases, the enzymes that work to break down cellulose. The more crystalline the structure, the stronger the fibers are. Fungal enzymes tested to date cannot easily break down fibers with high crystalline content so that the material can be converted into a biofuel. CelA, however, is agnostic to the level of crystalline content.

"CelA is able to break down cellulose with high crystallinity the same as low crystallinity, which has never been shown for any other cellulase," said Yannick Bomble, a senior research scientist at NREL and the senior author of the paper. "The better the cellulase is, the quicker you can convert biomass to simple sugars so the cheaper the process will get."

Bomble's co-authors from NREL are Roman Brunecky, Bryon Donohoe, John Yarbrough, Ashutosh Mittal, Larry Taylor, Daehwan Chung, and Michael Himmel. Other co-authors were Brian Scott, Hanshu Ding, Sarah Teter from Novozymes, and Jordan Russell and Janet Westpheling from the University of Georgia. Their research looked at how CelA performed in breaking down and interacting with the components of cell walls in [corn stover](#): glucan, xylan, and [lignin](#). Chemical pretreatments

were used on corn stover and silky fibers called cotton linters, leaving behind various amounts of the components and varying degrees of crystallinity, respectively. The experiments showed the degree of crystallinity didn't affect how well the enzyme performed.

The sticking point came when CelA encountered lignin, the component that provides rigidity to cell walls. With some pretreatment conditions, some lignin remained, and that stopped the enzyme. "If it binds to lignin, it's just stuck. It can't process or break down biomass any longer," Bomble said. "When that happens, you lose the enzyme. The more enzymes you lose to non-productive binding, the less efficient the conversion. That's usually the problem. This is why we're working on strategies to prevent CelA's binding to lignin but retain the vital affinity to cellulose."

**More information:** Roman Brunecky et al. The Multi Domain Caldicellulosiruptor bescii CelA Cellulase Excels at the Hydrolysis of Crystalline Cellulose, *Scientific Reports* (2017). [DOI: 10.1038/s41598-017-08985-w](https://doi.org/10.1038/s41598-017-08985-w)

Provided by National Renewable Energy Laboratory

Citation: Enzyme's worth to biofuels shown in latest research (2017, September 18) retrieved 13 July 2024 from <https://phys.org/news/2017-09-enzyme-worth-biofuels-shown-latest.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.