

An important process fueling harmful algal blooms investigated in Canadian water bodies

September 13 2017

For many Canadians, summer time means time at the lake, swimming, fishing, boating, and relaxing. Nothing can spoil this experience like bluegreen mats of muck, caused by algal blooms. These blooms negatively affect not only recreational activities but also put drinking water source, property values, wildlife, and human health at risk. In the 1970s, scientists discovered that the nutrient phosphorus caused algal blooms, which led to new regulations and improved sewage treatment. Nevertheless, blooms continue to plague many Canadian lakes. To investigate what might be happening, scientists looked to see whether phosphorus might be recirculating from the mud at the bottom of lakes back into the water.

In a new scientific study published today, scientists investigated an important, but poorly understood, process in Canada's aquatic ecosystems: the recycling of the algal nutrient, <u>phosphorus</u>, between mud at the bottom of lakes and the overlying water. This process can contribute to the formation of the <u>harmful algal blooms</u> that plague many lakes. Across the country tens or even hundreds of millions of dollars are invested every year to manage nutrient pollution but in some lakes legacy effects from nutrients deposited years ago can linger and delay recovery.

"Our main goal was to better understand where, when, and why this process occurs in Canadian fresh waters so that we can make improvements to how <u>algal blooms</u> are managed and develop realistic



goals for lake restoration," explained lead author Diane Orihel, an assistant professor at Queen's University. She added, "we need to stop dumping phosphorus into our lakes, because it's not only causing problems right now, but in many lakes, it continues to deteriorate water quality for our children and grandchildren."

Jason Venkiteswaran, an assistant professor at Wilfrid Laurier University pointed out that "We rely on phosphorus to fertilize our soils and produce the food the world needs. However, we don't know how long that phosphorus sticks around in our lakes after we allow it to wash off the land and down our drains. Our work here cautions that we should have different expectations for the recovery of different types of lakes across the country."

By critically reviewing data from 70 <u>water</u> bodies, the authors found that phosphorus release from sediments is a common phenomenon in Canadian fresh waters, but that rates of this process varied dramatically from lake to lake. "The highest rates of release were found in small prairie lakes in Manitoba, Saskatchewan, and Alberta, while the lowest rates were found in Canadian Shield lakes in Ontario and the Maritimes," said Nora Casson, assistant professor at the University of Winnipeg.

The authors of this study also identified the key factors controlling this process, such as oxygen, pH, geology and lake nutrient status—which often acts against the best efforts of <u>lake</u> managers—as well as identified areas where we know strikingly little. Helen Baulch, assistant professor at the University of Saskatchewan noted that "There's a lot more we need to know about internal phosphorus loading, such as if this process is important in the thousands of reservoirs of our country, and we have huge data gaps for this process in our northern lakes that are undergoing rapid change."



Featured in this article were case studies of Lake Simcoe, Lake Winnipeg, Lake of the Woods, Lake Erie, Lake Champlain, Cootes Paradise, and Lake Diefenbaker.

The journal article, "Internal phosphorus loading in Canadian fresh waters: a critical review and data analysis" by Diane Orihel (Queen's University), Helen Baulch (University of Saskatchewan), Nora Casson (University of Winnipeg), Rebecca North (University of Missouri), Chris Parsons (University of Waterloo), Dalila Seckar (Queen's University), and Jason Venkiteswaran (Wilfrid Laurier University) was published online today in the *Canadian Journal of Fisheries and Aquatic Sciences*. This article was selected by the journal as an "Editors' Choice" paper for 2017, which highlights articles of particularly high caliber and topical importance.

More information: Diane M. Orihel et al, Internal phosphorus loading in Canadian fresh waters: a critical review and data analysis, *Canadian Journal of Fisheries and Aquatic Sciences* (2017). DOI: 10.1139/cjfas-2016-0500

Provided by Canadian Science Publishing (NRC Research Press)

Citation: An important process fueling harmful algal blooms investigated in Canadian water bodies (2017, September 13) retrieved 3 May 2024 from <u>https://phys.org/news/2017-09-important-fueling-algal-blooms-canadian.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.