

Moth found to secrete distinct defensive fluids to ward off different types of predators

September 27 2017, by Bob Yirka



Wood tiger moth. Credit: Wikipedia

(Phys.org)—A team of researchers from the University of Jyväskylä in Finland and Technische Universität Braunschweig in Germany has found that the wood tiger moth secretes two different types of fluids from

different parts of its body to ward off two different types of predators. In their paper published in *Proceedings of the Royal Society B*, the group describes how they studied the moths in their lab and what they learned about them.

Wood tiger moths have coloring reminiscent of the tiger and are found in many parts of the world. In this new effort, the researchers took a closer look at how they ward off predators by collecting multiple specimens and bringing them back to the lab for study.

Prior research had shown that the moths excreted fluid from glands on the backs of their necks—a good location for warding off [birds](#), their main [predator](#). Birds eat moths and catch them by grabbing them by the back of the neck. Anecdotal evidence had suggested that when birds grab a wood tiger [moth](#), they tend to let it go, and have been seen trying to clean their beaks afterwards. Other anecdotal evidence also suggested that ants tend to avoid the moths, likely due to fluid excreted from the moth's anus.

The team started by taking a closer look at the fluid the moths produced from their necks and found that it contained chemical compounds known as pyrazines. The team treated oat flakes with the fluid and fed them to Eurasian blue tits and found that the birds were reluctant to eat them and became averse to them afterwards. The birds willingly ate flakes treated with the anal fluid and [sugar water](#).

Next, the team placed drops of sweetened fluid in disks laid in the path of ants—some of the disks also had chemicals taken from the moths. They report that the ants actually preferred the disks with neck fluid over just the sugar water, but clearly avoided those with anal fluid.

Based on their findings, the researchers conclude that the moths produce different [fluid](#) deterrents to ward off different types of predators.

More information: Bibiana Rojas et al. How to fight multiple enemies: target-specific chemical defences in an aposematic moth, *Proceedings of the Royal Society B: Biological Sciences* (2017). [DOI: 10.1098/rspb.2017.1424](https://doi.org/10.1098/rspb.2017.1424)

Abstract

Animals have evolved different defensive strategies to survive predation, among which chemical defences are particularly widespread and diverse. Here we investigate the function of chemical defence diversity, hypothesizing that such diversity has evolved as a response to multiple enemies. The aposematic wood tiger moth (*Arctia plantaginis*) displays conspicuous hindwing coloration and secretes distinct defensive fluids from its thoracic glands and abdomen. We presented the two defensive fluids from laboratory-reared moths to two biologically relevant predators, birds and ants, and measured their reaction in controlled bioassays (no information on colour was provided). We found that defensive fluids are target-specific: thoracic fluids, and particularly 2-sec-butyl-3-methoxypyrazine, which they contain, deterred birds, but caused no aversive response in ants. By contrast, abdominal fluids were particularly deterrent to ants, while birds did not find them repellent. Our study, to our knowledge, is the first to show evidence of a single species producing separate chemical defences targeted to different predator types, highlighting the importance of taking into account complex predator communities in studies on the evolution of prey defence diversity.

© 2017 Phys.org

Citation: Moth found to secrete distinct defensive fluids to ward off different types of predators (2017, September 27) retrieved 18 April 2024 from <https://phys.org/news/2017-09-moth-secrete-distinct-defensive-fluids.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.