

Study identifies likely scenarios for global spread of devastating crop disease

September 25 2017



Wheat infected with stem rust. Credit: CIMMYT

Stem rust, named for the blackening pustules that infect plant stems, caused devastating crop epidemics and famine for centuries before being tamed by fungicides and resistance genes.

Since the turn of the century, however, aggressive new strains have emerged—such as 'Ug99', first detected in Uganda in 1999—that infect widely grown varieties of wheat. These diseases threaten to disperse trillions of pathogenic fungal spores on winds across countries and continents. The current global economic loss from wheat [stem rust](#) is approximately U.S. \$1 billion a year.

The fear is that these airborne and highly virulent strains could spread from known sites to some of the world's most important 'breadbasket' regions, such as the Punjab in South Asia, where these strains have not yet been detected.


Now, a team of scientists of the University of Cambridge, the UK Met Office and CIMMYT (International Maize and Wheat Improvement Centre) have adapted modelling systems previously used to forecast, ash dispersal from erupting volcanoes and radiation from nuclear accidents (NAME), to predict when and how Ug99 and other such strains are most likely to spread.

The research, published today in the journal *Nature Plants*, quantifies for the first time the circumstances—routes, timings and outbreak sizes—under which dangerous strains of stem rust pose a threat from long-distance dispersal out of East Africa to the large wheat-producing

areas in India and Pakistan.

The results highlight the role of Yemen as a potential 'stepping stone' for the transmission of the [disease](#) between continents. The key scenario for disease spread is from Yemen directly to Pakistan or India. In case of a large outbreak in Eastern Yemen results indicate a 30% chance for transmission to occur.

Another important scenario for wheat rust spread is from Yemen through Middle Eastern countries, in particular Iran, to Central and South Asia. If Iran were to suffer a moderate outbreak of Ug99—more than 1000 hectares—then spores would likely spread to Afghanistan, and from there potentially further to the northern plains of Pakistan and India. However, transmission along this route is restricted to a relatively short time-window in March and April, before wheat is typically harvested in South Asia.



(A) This is a long-distance dispersal network of spores between all major wheat producing countries in Southern/East Africa, the Middle East and Central/South Asia. Nodes represent countries; communities of the same colour indicate regions with high airborne connectivity; the size of nodes indicates node-strength; pie charts show the fraction of out-strength to total node strength (indicating donor and receptor countries). (B) Spore transmission frequencies along principal migration routes in the Rift Valley zone for the scenario of large outbreaks. Credit: Matthew Hort

"New races of wheat rust are threatening wheat worldwide, and we need to know which areas are at risk," said senior author Prof Chris Gilligan, from Cambridge's Department of Plant Sciences.

"From our work, we now believe that if we start to see Ug99 or other new wheat rust strains take hold in Yemen in early spring then action must be taken immediately to mitigate the risk of further spread."

However, the modelling work also offers some encouraging news: the airborne transmission of the disease from East African countries directly to South Asia is highly unlikely, with transmission events possible only on less than one day a year.

The scientific team used field disease surveys from the International Maize and Wheat Improvement Centre and weather data from the UK Met Office as key input for the modelling framework.

"This research has allowed us to obtain the first quantitative estimates of long-term airborne spore transmission frequencies for different outbreak scenarios. We compiled risk assessments for pathogen dispersal from key disease locations to important wheat-producing countries. These assessments can effectively inform surveillance and control strategies,"

said Cambridge's Marcel Meyer, the study's first author.

The team say their work, including 3-D spore dispersal animations and a catalogue of dispersal trends (indicating likely directions, frequencies, pathogen loads), provides new ways to raise awareness, communicate risks, and inform agricultural stakeholders.

Their modelling framework can be applied as a tool to analyse risks in case new disease strains should be uncovered in other geographic areas. This has already helped in estimating dispersal risks from sites of other diseases in Europe and Siberia. In ongoing work the team is developing an Early Warning System forecasting rust risk in Ethiopia, East Africa's largest [wheat](#) producing country.

"The combined expertise from plant sciences and atmospheric dispersion sciences has delivered ground breaking tools that highlight the risks, and support the management of the devastating potential of these diseases," said Dr Matthew Hort, co-author from the UK's Met Office.

More information: M. Meyer et al, Quantifying airborne dispersal routes of pathogens over continents to safeguard global wheat supply, *Nature Plants* (2017). [DOI: 10.1038/s41477-017-0017-5](https://doi.org/10.1038/s41477-017-0017-5)

Provided by University of Cambridge

Citation: Study identifies likely scenarios for global spread of devastating crop disease (2017, September 25) retrieved 4 July 2024 from <https://phys.org/news/2017-09-scenarios-global-devastating-crop-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.