

New statistical approach for environmental measurements lets the data determine how to model extreme events

September 12 2017

Modeling environmental data, such as regional wind speed or temperature, is a complicated business. To model data statistically requires significant assumptions about its behavior over time and space—yet arriving at those assumptions requires an understanding of the data that can generally only be obtained by modeling. It's a catch-22 that presents a major obstacle to progress in large-scale environmental and climate modeling, particularly for extreme events.

KAUST researcher Raphaël Huser, in collaboration with colleagues from France and Switzerland, has now developed a modeling framework that lets the data define its own behavior around extreme events without the need for restrictive predetermined assumptions.

"Environmental extremes, such as extreme [wind](#) gusts, floods, or heat waves, are often spatially dependent," explains Huser. "That is, two neighboring measurement stations may, and often do, experience extreme events simultaneously. But does this dependence stabilize or weaken as the event becomes more extreme? Classical statistical models require the nature of this extremal dependence to be defined before modeling, but because extreme events are scarce, it can be very difficult if not impossible to correctly guess the dependence class in advance."

Classical statistical models that account for extreme events are known as asymptotic models. The choice of asymptotic extremal dependence type

determines how the model extrapolates to events even more extreme than those present in the data. It comes bundled with other implied assumptions that are not always realistic environmentally, with the result that such models may incorrectly assess the likelihood of future extreme events.

"We developed a suite of flexible geostatistical 'subasymptotic' models using a general Gaussian basis that captures both types of asymptotic dependence," says Huser. "Our models are more flexible and easier to use, especially for higher-dimensional data collected at many monitoring stations."

Through simulations of measured wind speeds, Huser's team showed that their Gaussian-scale mixtures model can accurately estimate the extremal dependence type. It also outperforms other typical models over a range of performance metrics with a good fit to the data and more realistic spatial prediction of extreme wind speeds at unobserved locations.

"The most important result of our work is that we no longer need to fix the asymptotic dependence class in advance but can let the data speak for itself," says Huser. "This [model](#) is applicable to a wide range of [environmental data](#) and will help improve our modeling and prediction of [extreme events](#)."

More information: Raphaël Huser et al, Bridging asymptotic independence and dependence in spatial extremes using Gaussian scale mixtures, *Spatial Statistics* (2017). [DOI: 10.1016/j.spasta.2017.06.004](https://doi.org/10.1016/j.spasta.2017.06.004)

Provided by King Abdullah University of Science and Technology

Citation: New statistical approach for environmental measurements lets the data determine how to model extreme events (2017, September 12) retrieved 25 April 2024 from <https://phys.org/news/2017-09-statistical-approach-environmental-extreme-events.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.