


## Ultrafast snapshots of relaxing electrons in solids

September 16 2017





Attosecond flashes of light and x-rays take snapshots of fleeting electrons in



solids. Credit: MPQ, Attoelectronics Group

When x-rays shine onto solid materials or large molecules, an electron is pushed away from its original place near the nucleus of the atom, leaving a hole behind. For a long time, scientists have suspected that the liberated electron and the positively charged hole form a new kind of quasiparticle—known as 'core-exciton'. But so far, there has not yet been a real proof of its existence. Scientists have a wide range of tools to track excitons in semiconductors in real-time. Those are generated by ordinary light, and can be employed in various applications in optoelectronics and microelectronics. On the contrary, core-excitons are extremely shortlived, and up to now, no technique was available to track their motion and deduce their properties.

"Core-excitons live for a very short time because their interactions with other particles in the solid quickly stops their motion," said Antoine Moulet, leading author in this work. "In quantum mechanics we say that the exciton loses its coherence," he adds.

A key tool to track the dynamics of core-excitons has been the development of attosecond light flashes in the optical range. The work



was published by the Attoelectronics group last year.

"In our experiment we use <u>x-ray flashes</u> to light up core-excitons in solids, whereas the optical attosecond pulses provide the possibility to resolve this motion in real-time," says Julien Bertrand, a former researcher in the group of Goulielmakis, at present assistant professor at Laval University, Canada. "The combination of both allowed us to take snapshots of the motion of core-excitons which lived for approximately 750 attoseconds."

But the study was not limited to capturing these fleeting motions inside solids. "We were able to acquire quantitative information about the properties of core-excitons such as their miniature dimension which were merely bigger than that of a single atom, or how easily they are polarized by visible light," says Goulielmakis. "Our technique advances excitonics, i.e. the measurement, the control and the application of excitons in the x-ray regime. But at the same time, it is a general tool for studying ultrafast x-ray initiated processes in solids on their natural time scales. Such a capability has never before been possible in x-ray science."

The team now envisages applications of their technique for studying ultrafast processes at interfaces of solids, and new routes to realize ultrafast switches for x-ray radiation based on optical light fields. "With x-ray free electron lasers rapidly proliferating around the world, the capability of controlling x-rays with visible light becomes increasingly important," says Goulielmakis.

**More information:** A. Moulet et al. Soft x-ray excitonics, *Science* (2017). <u>DOI: 10.1126/science.aan4737</u>



Citation: Ultrafast snapshots of relaxing electrons in solids (2017, September 16) retrieved 3 May 2024 from <u>https://phys.org/news/2017-09-ultrafast-snapshots-electrons-solids.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.