

# Study indicates arsenic can cause cancer decades after exposure ends

October 24 2017

---



A large sample of native arsenic. Credit: Aram Dulyan/Public Domain

A new paper published in the *Journal of the National Cancer Institute*

shows that arsenic in drinking water may have one of the longest dormancy periods of any carcinogen. By tracking the mortality rates of people exposed to arsenic-contaminated drinking water in a region in Chile, the researchers provide evidence of increases in lung, bladder, and kidney cancer even 40 years after high arsenic exposures ended.

Inorganic arsenic is naturally present at high levels in the groundwater of many countries, creating an important public health issue affecting millions of people. Severe health effects have been observed in populations drinking arsenic-contaminated water over long periods, and research has established that [drinking water](#) contaminated with arsenic causes skin cancer and several internal cancers such as lung, bladder, and [kidney cancer](#), as well as cardiovascular disease and other adverse outcomes.

The water source in Antofagasta, a city in northern Chile, experienced a sudden major increase in arsenic water concentrations in 1958, followed by a major reduction in exposure when an arsenic removal plant was installed in 1970. As the driest inhabited place on earth, everyone living in Antofagasta within this time period had to drink from city water sources with known [arsenic concentrations](#), thus exposing inhabitants to high concentrations of arsenic. Identifying a clear relationship between arsenic exposure and cancer [mortality rates](#), the study found that lung, bladder, and kidney [cancer mortality rates](#) started to increase about 10 years after the high exposures commenced and did not peak until at least 20 years after exposure reduction began. For both men and women, mortality rates for these types of cancer remained high up to 40 years after the highest exposure stopped.

Although the researchers plan to continue studying this population, they can already conclude that the delay between exposure to arsenic and the development of related cancers could be one of the longest of any human carcinogen. These findings not only add important scientific information

on latency patterns, they also may have direct public health implications.

The long latency after exposure reduction means the incidence of arsenic-related diseases is likely to remain very high for many years after arsenic exposures have stopped. Possible long-term interventions to reduce mortality and morbidity after high exposures end include disease screening, reducing important co-exposures, treatment and health services resource planning, and increasing public awareness of arsenic health effects.

**More information:** Allan H. Smith et al, Lung, Bladder, and Kidney Cancer Mortality 40 Years After Arsenic Exposure Reduction, *JNCI: Journal of the National Cancer Institute* (2017). [DOI: 10.1093/jnci/djx201](https://doi.org/10.1093/jnci/djx201)

Provided by Oxford University Press

Citation: Study indicates arsenic can cause cancer decades after exposure ends (2017, October 24) retrieved 25 April 2024 from <https://phys.org/news/2017-10-arsenic-cancer-decades-exposure.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--