

# Carbon coating gives biochar its garden-greening power

October 20 2017

---



Field experiment in Switzerland, with setup of compost windrows from mixed manure before adding the biochar. Credit: Nikolas Hagemann/University of Tübingen

For more than 100 years, biochar, a carbon-rich, charcoal-like substance made from oxygen-deprived plant or other organic matter, has both


delighted and puzzled scientists. As a soil additive, biochar can store carbon and thus reduce greenhouse gas emissions, and it can slow-release nutrients to act as a non-toxic fertilizer.

But the precise chemistry by which biochar stores nutrients and promotes plant growth has remained a mystery, so its commercial potential has been severely limited.

Now, an international team of researchers, with key contributions by Colorado State University experts, has illuminated unprecedented detail and mechanistic understanding of biochar's seemingly miraculous properties. The *Nature Communications* study, led by Germany's University of Tuebingen and published Oct. 20, demonstrated how composting of biochar creates a very thin organic [coating](#) that significantly improves the biochar's fertilizing capabilities. A combination of advanced analytical techniques confirmed that the coating strengthens the biochar's interactions with water and its ability to store soil nitrates and other nutrients.

This improved understanding of biochar's properties could trigger more widespread commercialization of biochar fertilizers. Such a change could reduce global dependence on inorganic nitrogen fertilizers that have served as modern food-production workhorses for more than a century.

The international collaboration included CSU's Thomas Borch, professor in the Department of Soil and Crop Sciences with joint appointments in chemistry and civil and environmental engineering, and research scientist Robert Young. The CSU team contributed high-resolution [mass spectrometry](#) performed at Florida State University's National High Magnetic Field Laboratory. Their data helped to confirm the composition of the biochar's nanoscale carbon coating.



The study analyzed the effects of a thin carbon coating on composted biochar.  
 Credit: Mihaela Albu, Austrian Cooperative Research, Graz; Wolfgang Gerber/Nikolas Hagemann, University of Tübingen

"To characterize a super-thin carbon coating on a carbon substrate is

nearly impossible," Borch said. "Our international team used many different advanced techniques to perform the analyses. Robert Young led our group's contribution of ultra-high resolution mass spectrometry to investigate the coating and probe its elemental makeup."

The study was led by Andreas Kappler, of the Center for Applied Geoscience at the University of Tuebingen, and geo-ecologist Nikola Hagemann. The authors set out to investigate biochar before and after composting with mixed manure. Using a combination of microscopic and spectroscopic analyses, the researchers found that dissolved organic substances played a key role in the composting of biochar and created the thin organic coating.

"This organic coating makes the difference between fresh and composted biochar," Kappler said. "The coating improves the biochar's properties of storing nutrients and forming further organic soil substances." Hagemann added that the coating also developed when untreated biochar was introduced into the soil - only much more slowly. Composting experiments were carried out on a small commercial scale using infrastructure and expertise of the Ithaka Institute in Switzerland.

Excessive use of mineral nitrogen fertilizers or liquid manure in agriculture has serious impacts on the environment. Such fertilizers cause the emission of nitrous oxide and result in nitrates leaching into the groundwater. As an eco-friendly alternative, scientists have suggested adding biochar as a [nutrient](#) carrier into the soil. But use of biochar on a large scale has not been economically viable because so little was known about exactly how it stores and releases nitrates.

"In agricultural crop production, higher yields usually only occur when biochar is applied together with nutrients from non-charred biomass such as liquid manure," Hagemann said. "Using [biochar](#) without adding nutrients or with pure mineral nutrients has proved to be far less

successful in many experiments."

**More information:** Nikolas Hagemann et al, Organic coating on biochar explains its nutrient retention and stimulation of soil fertility, *Nature Communications* (2017). [DOI: 10.1038/s41467-017-01123-0](https://doi.org/10.1038/s41467-017-01123-0)

Provided by Colorado State University

Citation: Carbon coating gives biochar its garden-greening power (2017, October 20) retrieved 17 April 2024 from

<https://phys.org/news/2017-10-carbon-coating-biochar-garden-greening-power.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--