

A new way to harness wasted methane

October 17 2017, by David L. Chandler



MIT chemistry professor Yogesh Surendranath and three colleagues have found a way to use electricity, which could potentially come from renewable sources, to convert methane into derivatives of methanol. The researchers developed a lowtemperature electrochemical process that would continuously replenish a catalyst material that can rapidly carry out the conversion. Credit: Massachusetts Institute of Technology



Methane gas, a vast natural resource, is often disposed of through burning, but new research by scientists at MIT could make it easier to capture this gas for use as fuel or a chemical feedstock.

Many oil wells burn off <u>methane</u>—the largest component of natural gas—in a <u>process</u> called flaring, which currently wastes 150 billion cubic meters of the gas each year and generates a staggering 400 million tons of carbon dioxide, making this process a significant contributor to global warming. Letting the gas escape unburned would lead to even greater environmental harm, however, because methane is an even more <u>potent</u> <u>greenhouse gas</u> than carbon dioxide is.

Why is all this methane being wasted, when at the same time natural gas is touted as an important "bridge" fuel as the world steers away from fossil fuels, and is the centerpiece of the so-called shale-gas revolution? The answer, as the saying goes in the real estate business, is simple: location, location.

The wells where methane is flared away are primarily being exploited for their petroleum; the methane is simply a byproduct. In places where it is convenient to do so, methane is captured and used to generate electrical power or produce chemicals. However, special equipment is needed to cool and pressurize <u>methane gas</u>, and special pressurized containers or pipelines are needed to transport it. In many places, such as offshore oil platforms or remote oil fields far from the needed infrastructure, that's just not economically viable.

But now, MIT chemistry professor Yogesh Surendranath and three colleagues have found a way to use electricity, which could potentially come from renewable sources, to convert methane into derivatives of methanol, a liquid that can be made into automotive fuel or used as a precursor to a variety of chemical products. This new method may allow for lower-cost methane conversion at remote sites. The findings,



described in the journal *ACS Central Science*, could pave the way to making use of a significant methane supply that is otherwise totally wasted.

"This finding opens the doors for a new paradigm of methane conversion chemistry," says Jillian Dempsey, an assistant professor of chemistry at the University of North Carolina, who was not involved in this work.

Existing industrial processes for converting methane to liquid intermediate chemical forms requires very high operating temperatures and large, capital-intensive equipment. Instead, the researchers have developed a low-temperature electrochemical process that would continuously replenish a catalyst material that can rapidly carry out the conversion. This technology could potentially lead to "a relatively lowcost, on-site addition to existing wellhead operations," says Surendranath, who is the Paul M. Cook Career Development Assistant Professor in MIT's Department of Chemistry.

The electricity to power such systems could come from wind turbines or solar panels close to the site, he says. This electrochemical process, he says, could provide a way to do the methane conversion—a process also known as functionalizing—"remotely, where a lot of the 'stranded' methane reserves are."

Already, he says, "methane is playing a key role as a transition fuel." But the amount of this valuable fuel that is now just flared away, he says, "is pretty staggering." That vast amount of wasted natural gas can even be seen in satellite images of the Earth at night, in areas such as the Bakken oil fields in North Dakota that light up as brightly as big metropolitan areas due to flaring. Based on World Bank estimates, global flaring of methane wastes an amount equivalent to approximately one-fifth of U.S. <u>natural gas</u> consumption.



When that gas gets flared off rather than directly released, Surendranath says, "you're reducing the environmental harm, but you're also wasting the energy." Finding a way to do methane conversion at sufficiently low cost to make it practical for remote sites "has been a grand challenge in chemistry for decades," he says. What makes methane conversion so tough is that the carbon-hydrogen bonds in the methane molecule resist being broken, and at the same time there's a risk of overdoing the reaction and ending up with a runaway process that destroys the desired end-product.

Catalysts that could do the job have been studied for many years, but they typically require harsh chemical agents that limit the speed of the reaction, he says. The key new advance was adding an electrical driving force that could be tuned precisely to generate more potent catalysts with very high reaction rates. "Since we're using electricity to drive the process, this opens up new opportunities for making the process more rapid, selective, and portable than existing methods," Surendranath says. And in addition, "we can access catalysts that no one has observed before, because we're generating them in a new way."

The result of the reaction is a pair of liquid chemicals, methyl bisulfate and methanesulfonic acid, which can be further processed to make liquid methanol, a valuable chemical intermediate to fuels, plastics, and pharmaceuticals. The additional processing steps needed to make methanol remain very challenging and must be perfected before this technology can be implemented on an industrial scale. The researchers are actively refining their method to tackle these technological hurdles.

"This work really stands out because it not only reports a new system for selective catalytic functionalization of methane to methanol precursors, but it includes detailed insight into how the system is able to carry out this selective chemistry. The mechanistic information will be instrumental in translating this exciting discovery into an industrial



technology," Dempsey says.

More information: Matthew E. O'Reilly et al. Catalytic Methane Monofunctionalization by an Electrogenerated High-Valent Pd Intermediate, *ACS Central Science* (2017). <u>DOI:</u> <u>10.1021/acscentsci.7b00342</u>

This story is republished courtesy of MIT News (web.mit.edu/newsoffice/), a popular site that covers news about MIT research, innovation and teaching.

Provided by Massachusetts Institute of Technology

Citation: A new way to harness wasted methane (2017, October 17) retrieved 2 May 2024 from <u>https://phys.org/news/2017-10-harness-methane.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.