

Mold contamination in sea salts could potentially spoil food

October 3 2017, by Joe Schwartz



Credit: CC0 Public Domain

Like fine wines, sea salts are artisanal products that inspire talk of terroir, texture and provenance. Now there's evidence that they can also be sources of spoilage molds.



Research from Cornell University mycologist Kathie Hodge and doctoral candidate Megan Biango-Daniels reveals varying levels of mold contamination in commercial sea salts. Among those molds were important <u>food</u> spoilage molds like *Aspergillus* and *Penicillium*, and even some notorious producers of mycotoxins.

"This new finding contradicts the conventional wisdom that salts are sterile ingredients," said Biango-Daniels. The research stressed the importance of understanding the risk of using sea <u>salt</u> during food production.

Starting with seven different commercial salts, the researchers extracted living fungi and grew them in the lab for identification. The fungi discovered in the salt have the potential to spoil food when used as an ingredient and can introduce mycotoxins or allergens when consumed.

At the levels discovered in the study, about 1.7 spores per gram, you're not risking your health by sprinkling sea salt on food you are about to eat. But big problems may result when sea salts are used at home or industrially to make cured meats, fermented pickles or brined cheeses that mature over time - when molds introduced with sea salt can begin to grow and spoil food, maybe even rendering it toxic.

"Fungi can survive in surprisingly hostile places. They can't increase or grow in a container of sea salt - nothing can - but spores of some fungi survive quite happily there. Later they can wake up and make trouble in our food," said Hodge, associate professor in the Plant Pathology and Plant-Microbe Biology Section in Cornell's College of Agriculture and Life Sciences.

Fungi likely make their way into sea salts during their production from seawater or during storage and packaging. Sea salt production starts with seawater trapped in outdoor ponds called salterns. As the seawater



evaporates, <u>salt crystals</u> form and are scraped up from the ponds and dried. Fungi may remain from microbes that live in the salterns, or their spores may fall in from the air over weeks of evaporation. Once <u>fungi</u> are in the salt, they have the potential to grow again once they encounter a moister environment. The researchers said microbial safety standards for this currently unregulated ingredient are needed to prevent food spoilage.

More information: Megan N. Biango-Daniels et al, Sea salts as a potential source of food spoilage fungi, *Food Microbiology* (2017). DOI: <u>10.1016/j.fm.2017.07.020</u>

Provided by Cornell University

Citation: Mold contamination in sea salts could potentially spoil food (2017, October 3) retrieved 1 May 2024 from <u>https://phys.org/news/2017-10-mold-contamination-sea-salts-potentially.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.