


Research clarifies nuclear hormone receptor function in plants

October 11 2017



Model for the mechanistic action of MED25 in cooperating with both genetic and epigenetic regulators in regulating jasmonate-induced activation of MYC2.
Credit: IGDB

The striking capacity for plants to adapt their growth and development to an ever-changing environment is mediated by diverse plant hormones that regulate virtually every aspect of plant life. In the past 10 to 15 years, enormous progress has been made in elucidating the nature of plant hormone receptors.

An important conceptual advance in hormone biology is the finding that, in addition to conventional membrane-bound [hormone receptors](#), the main receptors of several plant hormones are localized in the nucleus and directly linked to hormone-regulated [gene transcription](#). Despite this progress, the physiological significance and action mechanism of nuclear hormone receptors (NRs) in the plant kingdom remain enigmatic.

Jasmonate is a lipid-derived plant hormone that regulates a wide range of plant immunity and adaptive growth functions. Sensing of jasmonate by the nuclear-localized F-box protein CORONATINE INSENSITIVE 1 (COI1) triggers genome-wide transcriptional changes that are largely regulated by the master transcription factor MYC2. However, it remains unclear how COI1 relays hormone-specific regulatory signals to the RNA polymerase II (Pol II) general transcriptional machinery and chromatin.

A new study led by LI Chuanyou from the Institute of Genetics and Developmental Biology of the Chinese Academy of Sciences has revealed that Mediator, an evolutionarily conserved multi-subunit coactivator complex whose activity is essential for Pol II-dependent gene transcription, directly links COI1 to Pol II and chromatin during jasmonate signaling.

They found that, in the resting stage, the Mediator subunit MED25 brings COI1 to MYC2 target promoters through physical interaction; upon hormone elicitation, MED25 facilitates COI1-dependent degradation of JAZ transcriptional repressors, which favors

MYC2-directed transcription of jasmonate-responsive genes.

In addition, they found that MED25 also physically and functionally interacts with HISTONE ACETYLTRANSFERASE1 (HAC1), an evolutionarily conserved histone modification enzyme that selectively regulates histone (H) 3 lysine (K) 9 acetylation (H3K9ac) of MYC2 target promoters during jasmonate signaling. Moreover, MED25 cooperates with both COI1 and HAC1 on MYC2 target promoters.

These new observations, together with their previous finding that MED25 bridges the master transcription factor MYC2 and Pol II for pre-initiation complex (PIC) assembly during jasmonate-regulated gene transcription (Chen et al., 2012, The Plant Cell), reveal that MED25 acts as a master coordinator to integrate the actions of both genetic and epigenetic regulators into a concerted transcriptional program.

The signaling paradigm whereby Mediator links hormone receptors to transcriptionally active chromatin likely operates in other plant hormones whose receptors are localized in the nucleus.

Moreover, the researchers showed that the overall protein domain composition of plant MED25 is largely similar to that of its animal counterpart, which engages in a ligand-dependent interaction with retinoic acid receptor (RAR) and several other NRs.

In the context that animal Mediator was first biochemically isolated as a thyroid [hormone](#) receptor-associated protein (TRAP) complex, and has been shown to be an indispensable NR-interacting coactivator, this study reveals a scenario in which [plants](#) and animals have evolved distinct, but nonetheless largely similar, mechanisms for NR activation at the level of transcriptional regulation.

Thus, this investigation lays an important foundation not only for

investigating the action of Mediator in other [plant hormones](#), but also for examining the similarities and differences between plant and animal NR systems.

More information: Chunpeng An et al, Mediator subunit MED25 links the jasmonate receptor to transcriptionally active chromatin, *Proceedings of the National Academy of Sciences* (2017). [DOI: 10.1073/pnas.1710885114](#)

Provided by Chinese Academy of Sciences

Citation: Research clarifies nuclear hormone receptor function in plants (2017, October 11)
retrieved 3 May 2024 from
<https://phys.org/news/2017-10-nuclear-hormone-receptor-function.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--