

# Pheasant roadkill peaks in autumn and late winter

October 3 2017

---



Pheasants are commonly roadkill, but the recent increase in their artificial rearing and release has changed the time of year when most fatalities occur  
Credit: Bev Milne

Chickens' motives for crossing the road are often questioned - but

pheasants should probably avoid it altogether, new research suggests. Researchers from the universities of Exeter and Cardiff compared roadkill figures from the 1960s and 2010s - before and after the start of mass release programmes of pheasants for shooting - and found pheasants remain disproportionately likely to be run over compared to other birds. "There may be a number of reasons why pheasants are so commonly killed on the roads, including their short flight distances and relatively small brains," said Dr Joah Madden, of the University of Exeter.

"Our research shows that large-scale release of [pheasants](#) has not changed their likelihood of being killed, but it has changed the times of year when they are being killed."

The peak times of year for pheasant roadkill have changed from early summer in the 1960s to autumn and late winter now.

These twin peaks coincide with times in the year when captive-bred pheasants are released from pens, and when supplementary feeding ceases following the end of the shooting season.

"Pheasants in the 2010s are no longer susceptible during their June-August breeding season, unlike in the 1960s, perhaps because relatively few breed successfully," Dr Madden said.

"Instead, roadkill first peaks in September-November as pheasants disperse from release pens. It then declines over winter, but when supplementary feeding ceases in February, we see a second peak.

"Captive-bred pheasants may be at risk after release from pens because they have not learned survival skills.

"Being raised in the absence of parents, they simply lack the opportunity

to learn."





Pheasants are commonly roadkill, but the recent increase in their artificial rearing and release has changed the time of year when most fatalities occur

Credit: Bev Milne

Dr Madden, whose work was funded by the European Research Council, added: "Certain simple steps might help reduce pheasant roadkill, such as feeding them away from roads and continuing feeding them from February to May, after the shooting season has ended, to stop them wandering on to roads."

The researchers used roadkill figures from citizen science group Project Splatter, established and coordinated by Dr Sarah Perkins, of Cardiff University.

"Our work demonstrates how changes in animal behaviour can be revealed by [roadkill](#) data reported by members of the public, and the value of citizen science," Dr Perkins said.



Pheasant. Credit: Shaun Ferguson

**More information:** Why did the pheasant cross the road? Long term road mortality patterns in relation to management changes, *Royal Society Open Science*, [rsos.royalsocietypublishing.org ... /10.1098/rsos.170617](https://rsos.royalsocietypublishing.org/.../10.1098/rsos.170617)

Provided by University of Exeter

Citation: Pheasant roadkill peaks in autumn and late winter (2017, October 3) retrieved 30 April 2024 from <https://phys.org/news/2017-10-pheasant-roadkill-peaks-autumn-late.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.