


Physicists gain understanding of how bubbles at the edge of plasmas can drain heat and reduce fusion reaction efficiency

October 19 2017



PPPL physicist Michael Churchill. Credit: Elle Starkman

To fuse hydrogen atoms into helium, doughnut-shaped devices called tokamaks must maintain the heat of the ultrahot plasma they control. But like boiling water, plasma has blobs, or bubbles, that percolate within the plasma edge, reducing the performance of the plasma by taking away heat that sustains the fusion reactions.

Now, scientists at the U.S. Department of Energy's (DOE) Princeton Plasma Physics Laboratory (PPPL) have completed new simulations that could provide insight into how blobs at the plasma [edge](#) behave. The simulations, produced by a code called XGC1 developed by a national team based at PPPL, performed kinetic simulations of two different regions of the plasma edge simultaneously. This ability produces a more fundamental and fuller picture of how heat moves from plasma to the walls, potentially causing damage.

"In simulations, we often separate two areas at the plasma edge known as the pedestal and the scrape-off layer and focus on one or the other," said PPPL physicist Michael Churchill, lead author of a paper describing the results in the journal *Plasma Physics and Controlled Fusion*. "XGC1 is unique because it is able to simulate both regions simultaneously, using kinetic ion and electron equations. In fact, it is important to include both regions in simulations because they affect each other."

Simulations allow scientists to explore plasma, the fourth and hottest state of matter in which electrons are separated from atomic nuclei, without running [physical experiments](#) that could be costly. They also sometimes provide insights that physical experiments do not.

Simulations of turbulence at the edge of the plasma, near where the plasma approaches a tokamak's interior wall, are particularly important. The more that scientists understand such turbulence, the better able they will be to prevent moving blobs of plasma from forming in the plasma edge. If not controlled, these blobs could drain large amounts of heat from the confined plasma, and possibly either damage plasma-facing

components or hinder the [fusion reactions](#).

The XGC1 code simulated plasma in high-confinement mode, or H-mode, a set of conditions that helps plasma retain its heat. In H-mode, the results showed, a large number of blobs form between the pedestal and the scrape-off layer, two conditions near the edge, and move towards the outer edge, crossing the magnetic field lines as they go.

Blobs play an important role in the outward movement of particles in plasma. Blobs cause approximately 50 percent of the particle loss at the plasma edge, and researchers have observed blobs in a wide range of plasma devices, including tokamaks, figure-eight-shaped fusion devices known as stellarators, and linear machines. "The big picture is that blobs can pull energy and particles out of the [plasma](#), and you don't want that," Churchill said. "You want to keep things confined."

More information: R M Churchill et al, Pedestal and edge electrostatic turbulence characteristics from an XGC1 gyrokinetic simulation, *Plasma Physics and Controlled Fusion* (2017). [DOI: 10.1088/1361-6587/aa7c03](#)

Provided by Princeton Plasma Physics Laboratory

Citation: Physicists gain understanding of how bubbles at the edge of plasmas can drain heat and reduce fusion reaction efficiency (2017, October 19) retrieved 27 April 2024 from <https://phys.org/news/2017-10-physicists-gain-edge-plasmas-fusion.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
