

Scientists discover more about the ingredients for star formation

October 10 2017, by Luca Cortese



Comparison between the stellar (top) and molecular hydrogen (bottom) distribution in very gas-rich galaxies three billion years younger than the Milky Way. Optical data is from the Sloan Digital Survey whereas molecular hydrogen maps have been obtained using the Atacama Large Millimetre Array.

Astronomers have shed fresh light on the importance of hydrogen atoms in the birth of new stars.

Only hydrogen molecules are thought to directly fuel star formation but research published today shows there are more <u>hydrogen atoms</u> than molecules even in young galaxies that are making a lot of stars.



Astrophysicist Dr Luca Cortese, from The University of Western Australian node of the International Centre for Radio Astronomy Research, said <u>new stars</u> were constantly forming in the Universe.

"New stars are born in dense clouds of gas and dust that are found in most galaxies," he said.

"Our own Milky Way forms about one new star a year on average."

In the local Universe close to us about 70 per cent of the <u>hydrogen gas</u> is found in individual atoms, while the rest is in molecules.

Astronomers had expected that as they looked back in time, younger galaxies would contain more and more molecular hydrogen until it dominated the gas in the galaxy. Instead, they found that atomic hydrogen made up the majority of gas in younger galaxies too.

This is true even in galaxies under conditions similar to 'cosmic noon', a period about seven billion years after the Big Bang when the rate of star formation in the Universe reached its peak.

Dr Cortese said that in the last decade astronomers had discovered young, star-forming galaxies at cosmic noon with 10 times more <u>hydrogen molecules</u> than the Milky Way.

With such large reservoirs of molecular hydrogen, no room seemed to be left for a comparable amount of cold atomic gas. Unfortunately, it is currently impossible to detect hydrogen atoms at such large distances and verify this expectation.

Instead, Dr Cortese and his team discovered a population of galaxies three billion years younger than the Milky Way hosting gas reservoirs at least as large as those of galaxies at the cosmic noon.











Comparison between the stellar (top) and molecular hydrogen (bottom) distribution in a very gas-rich galaxy (AGC191728) that is three billion years younger than the Milky Way. Credit: ICRAR

"What we found is that despite hosting 10 billion solar masses of molecular gas these young galaxies turn out to be very, very rich in atomic hydrogen as well," Dr Cortese said.

"The balance between atomic and molecular hydrogen is pretty much the same as in the Milky Way. In other words, it's still dominated by atomic gas."

The research used data from two of the world's most powerful radio telescopes, the Arecibo Observatory in Puerto Rico and the European Southern Observatory's Atacama Large Millimeter/submillimeter Array in Chile.

ICRAR astrophysicist Dr Barbara Catinella, a co-author on the research, said the findings had tremendous implications for our understanding of the early Universe.

"It shows that we cannot neglect <u>atomic hydrogen</u> even in galaxies that contain tens of billions of solar masses of molecular <u>hydrogen</u>," she said.

"Only the advent of future radio telescopes such as the Square Kilometre Array will allow us to get a complete picture of the role of cold gas in the <u>star formation</u> cycle."

A further finding from the study was that the galaxies rich in <u>molecular</u>



hydrogen were not very turbulent.

Usually, these <u>galaxies</u> would be expected to be very turbulent to prevent the collapse of the gas into stars.

The research was published in the Astrophysical Journal Letters.

More information: ALMA shows that gas reservoirs of star-forming disks over the past 3 billion years are not predominantly molecular. *arXiv*. <u>arxiv.org/abs/1709.07933</u>

Provided by University of Western Australia

Citation: Scientists discover more about the ingredients for star formation (2017, October 10) retrieved 30 April 2024 from https://phys.org/news/2017-10-scientists-ingredients-star-formation.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.