

What soot-covered, hundred-year-old birds can tell us about saving the environment

October 9 2017



Horned Larks from The Field Museum's collections, with gray birds from the turn of the century and cleaner birds from more recent years when there was less soot in the atmosphere. Credit: (c) Carl Fuldner and Shane DuBay, The University of Chicago and The Field Museum

Horned Larks are cute little songbirds with white bellies and yellow chins—at least, now they are. A hundred years ago, at the height of urban smoke pollution in the US, their pale feathers were stained dark



gray by the soot in the atmosphere. A new paper in the *Proceedings of the National Academy of Sciences* shows that the discoloration of birds in museum collections can be used to trace the amount of black carbon in the air over time and the effects of environmental policy upon pollution.

"The soot on these birds' feathers allowed us to trace the amount of black carbon in the air over time, and we found that the air at the turn of the century was even more polluted than scientists previously thought," says Shane DuBay, a graduate student at The Field Museum and the University of Chicago and one of the authors of the study. He and coauthor Carl Fuldner, also a graduate student at UChicago, analyzed over a thousand birds collected over the last 135 years to determine and quantify the effects of soot in the air over cities in the Rust Belt.

"If you look at Chicago today, the skies are blue. But when you look at pictures of Beijing and Dehli, you get a sense for what US cities like Chicago and Pittsburgh were once like," says DuBay. "Using museum collections, we were able to reconstruct that history."

Ornithologists at The Field Museum have long known that bird specimens in the collection from the early 1900s were visibly darker than expected, and atmospheric soot was the suspect.

"When you touch these birds, you get traces of soot on your hands. We'd wear white gloves while handling them, and the gloves would come away stained, like when you get ink on your fingertips reading a newspaper," says DuBay. That's because the soot in the air clung to the birds' feathers like dust to a feather duster. "These birds were acting as air filters moving through the environment," adds DuBay.





Field Sparrows from The Field Museum's collections, with gray birds from the turn of the century and cleaner birds from more recent years when there was less soot in the atmosphere. Credit: (c) Carl Fuldner and Shane DuBay, The University of Chicago and The Field Museum

Birds were also ideal candidates for the study because they molt and grow a new set of feathers every year, meaning that the soot on them had only been accumulating for the past year when they were collected. And there was an apparent trend: old birds were dirtier, and new birds were cleaner.

To measure the changes in sootiness over the years, DuBay and Fuldner turned to a novel approach: photographing the birds and measuring the light reflected off of them. Fuldner, a photo historian who focuses on images of the environment, worked with DuBay to develop a method for analyzing the birds using photography. The birds photographed,



numbering over a thousand, were all from five species that breed in the Manufacturing Belt and have lots of white feathers that showed off the soot.

The images, depicting the contrast between the soiled gray birds and the clean white ones, are dramatic. "The photographs give the project a visceral dimension—you make a connection to the images," says Fuldner.

DuBay and Fuldner plotted the amount of light bouncing off the birds' feathers according to the year the birds were collected. To make sense of their findings, the pair of researchers then delved into the social history of urban air pollution.

"The changes in the birds reflect efforts, first at the city level but eventually growing into a national movement, to address the smoke problem," says Fuldner. "We are actually able to go back and see how effective certain policy approaches were."

"We were surprised by the precision we were able to achieve," says DuBay. "The soot on the birds closely tracks the use of coal over time. During the Great Depression, there's a sharp drop in black carbon on the birds because coal consumption dropped—once we saw that, it clicked." The amount of soot on the birds rebounded around World War II, when wartime manufacturing drove up coal use, and dropped off quickly after the war, around when people in the Rust Belt began heating their homes with natural gas piped in from the West rather than with coal.





Horned Larks from The Field Museum's collections, with gray bird from the turn of the century and cleaner bird from more recent years when there was less soot in the atmosphere. Credit: (c) Carl Fuldner and Shane DuBay, The University of Chicago and The Field Museum

"The fact that the more recent <u>birds</u> are cleaner doesn't mean we're in the clear," DuBay notes. "While the US releases far less black carbon into the atmosphere than we used to, we continue to pump lessconspicuous pollutants into our atmosphere—those pollutants just aren't as visible as soot. Plus, many people around the world still experience <u>soot</u>-choked air in their cities."

Analysis of atmospheric black carbon might assist scientists studying climate change. "We know black carbon is a powerful agent of climate change, and at the turn of the century, black carbon levels were worse than previously thought," says DuBay. "I hope that these results will help



climate and atmospheric scientists better understand the effects of black carbon on climate."

And for both DuBay and Fuldner, being able to apply their research beyond their respective fields of evolutionary biology and photographic history was both unexpected and rewarding.

"As a historian, one of the questions I always ask is, 'What is the point of this research to the way we live now?' In this case the answer quickly became clear," says Fuldner. "Filling in a blank space in the historical record of something as large as air pollution in American cities, and being able to share that with atmospheric scientists who study the effects of <u>black carbon</u> on the climate, is extraordinary."

"This study shows a tipping point when we moved away from burning dirty coal, and today, we're at a similar pivotal moment with fossil fuels," says DuBay. "In the middle of the 20th century, we made an investment in infrastructure and regulated fuel sources—hopefully, we can take that lesson and make a similar transition now to more sustainable, renewable energy sources that are more efficient and less harmful to our environment."

DuBay notes that in addition to the environmental implications of the project, their work also shows the importance of <u>museum collections</u> like those they used from The Field Museum in Chicago, the Carnegie Museum of Natural History in Pittsburgh, and the University of Michigan Museum of Zoology in Ann Arbor. "I hope this study exposes collections as a valuable resource to address present day environmental concerns," says DuBay. "This paper shows the ways that natural history collections can be used, underlining the value in collections and in continuing to build collections, to help us improve our understanding of human impacts on the natural world."



More information: Shane G. DuBay el al., "Bird specimens track 135 years of atmospheric black carbon and environmental policy," *PNAS* (2017). <u>www.pnas.org/cgi/doi/10.1073/pnas.1710239114</u>

Provided by Field Museum

Citation: What soot-covered, hundred-year-old birds can tell us about saving the environment (2017, October 9) retrieved 27 April 2024 from <u>https://phys.org/news/2017-10-soot-covered-hundred-year-old-birds-environment.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.