


# ALMA discovers infant stars surprisingly near galaxy's supermassive black hole

November 28 2017

---



An ALMA image of the center of the Milky Way galaxy showing the location of 11 young protostars within about 3 light-years of our galaxy's supermassive black hole. The lines indicate the direction of the bipolar lobes created by high-velocity jets from the protostars. The illustrated star in the middle of the image indicates the location of Sagittarius A\*, the 4 million solar mass supermassive black hole at the center of our galaxy. The next image is a zoom-in to one of the protostars. Credit: ALMA (ESO/NAOJ/NRAO), Yusef-Zadeh et al.; B. Saxton (NRAO/AUI/NSF)

At the center of our galaxy, in the immediate vicinity of its supermassive black hole, is a region wracked by powerful tidal forces and bathed in intense ultraviolet light and X-ray radiation. These harsh conditions, astronomers surmise, do not favor star formation, especially low-mass stars like our sun. Surprisingly, new observations from the Atacama Large Millimeter/submillimeter Array (ALMA) suggest otherwise.


ALMA has revealed the telltale signs of eleven [low-mass stars](#) forming perilously close—within three light-years—to the Milky Way's supermassive black hole, known to astronomers as Sagittarius A\* (Sgr A\*). At this distance, [tidal forces](#) driven by the supermassive black hole should be energetic enough to rip apart [clouds](#) of dust and gas before they can form stars.

The presence of these newly discovered protostars (the formative stage between a dense cloud of gas and a young, shining star) suggests that the conditions necessary to birth low-mass stars may exist even in one of the most turbulent regions of our galaxy and possibly in similar locales throughout the universe.

The results are published in the *Astrophysical Journal Letters*.

"Despite all odds, we see the best evidence yet that low-mass stars are

forming startlingly close to the [supermassive black hole](#) at the center of the Milky Way," said Farhad Yusef-Zadeh, an astronomer at Northwestern University in Evanston, Illinois, and lead author on the paper. "This is a genuinely surprising result and one that demonstrates just how robust [star formation](#) can be, even in the most unlikely of places."



Double-lobe feature produced by jets from one of the newly forming stars. ALMA discovered 11 of these telltale signs of star formation remarkably close

to the supermassive black hole at the center of our galaxy. Credit: ALMA (ESO/NAOJ/NRAO), Yusef-Zadeh et al.; B. Saxton (NRAO/AUI/NSF)

The ALMA data also suggest that these protostars are about 6,000 years old. "This is important because it is the earliest phase of star formation we have found in this highly hostile environment," Yusef-Zadeh said.

The team of researchers identified these protostars by seeing the classic "double lobes" of material that bracket each of them. These cosmic hourglass-like shapes signal the early stages of star formation. Molecules, like carbon monoxide (CO), in these lobes glow brightly in millimeter-wavelength light, which ALMA can observe with remarkable precision and sensitivity.


Protostars form from interstellar clouds of dust and gas. Dense pockets of material in these clouds collapse under their own gravity and grow by accumulating more and more star-forming gas from their parent clouds. A portion of this infalling material, however, never makes it onto the surface of the star. Instead, it is ejected as a pair of high-velocity jets from the protostar's north and south poles. Extremely turbulent environments can disrupt the normal procession of material onto a protostar, while intense radiation—from massive nearby stars and supermassive black holes—can blast away the parent cloud, thwarting the formation of all but the most massive of stars.

The Milky Way's galactic center, with its 4 million solar mass black hole, is located approximately 26,000 light-years from Earth in the direction of the constellation Sagittarius. Vast stores of interstellar dust obscure this region, hiding it from optical telescopes. Radio waves, including the millimeter and submillimeter light that ALMA sees, are able to penetrate this dust, giving radio astronomers a clearer picture of



the dynamics and content of this hostile environment.

Prior ALMA observations of the region surrounding Sgr A\* by Yusef-Zadeh and his team revealed multiple massive infant stars that are estimated to be about 6 million years old. These objects, known as proplyds, are common features in more placid star-forming regions, like the Orion Nebula. Though the galactic center is a challenging environment for star formation, it is possible for particularly dense cores of hydrogen gas to cross the necessary threshold and forge new stars.



Infant stars, like those recently identified near the supermassive black hole at the center of our galaxy, are surrounded by a swirling disk of dust and gas. In this artist's conception of an infant solar system, the young star pulls material from its surroundings into a rotating disk (right) and generates outflowing jets of material (left). Credit: Bill Saxton (NRAO/AUI/NSF)

The new ALMA observations, however, revealed something even more

remarkable, signs that eleven low-mass protostars are forming within 1 parsec – a scant 3 light-years – of the galaxy's central black hole. Yusef-Zadeh and his team used ALMA to confirm that the masses and momentum transfer rates – the ability of the protostar jets to plow through surrounding interstellar material – are consistent with young protostars found throughout the disk of our galaxy.

"This discovery provides evidence that star formation is taking place within clouds surprisingly close to Sagittarius A\*," said Al Wootten with the National Radio Astronomy Observatory in Charlottesville, Virginia, and co-author on the paper. "Though these conditions are far from ideal, we can envision several pathways for these stars to emerge."

For this to occur, outside forces would have to compress the gas clouds near the center of our galaxy to overcome the violent nature of the region and allow gravity to take over and form stars. The astronomers speculate that high-velocity gas clouds could aid in star formation as they force their way through the interstellar medium. It is also possible that jets from the black hole itself could be plowing into the surrounding gas clouds, compressing material and triggering this burst of star formation.

"The next step is to take a closer look to confirm that these newly formed stars are orbited by disks of dusty gas," concluded Mark Wardle, an astronomer at Macquarie University in Sydney, Australia, and co-investigator on the team. "If so, it's likely that planets will eventually form from this material, as is the case for young [stars](#) in the galactic disk."

**More information:** F. Yusef-Zadeh et al. ALMA Detection of Bipolar Outflows: Evidence for Low-mass Star Formation within 1 pc of Sgr A\*, *The Astrophysical Journal* (2017). [DOI: 10.3847/2041-8213/aa96a2](https://doi.org/10.3847/2041-8213/aa96a2)

Provided by National Radio Astronomy Observatory

Citation: ALMA discovers infant stars surprisingly near galaxy's supermassive black hole (2017, November 28) retrieved 23 April 2024 from <https://phys.org/news/2017-11-alma-infant-stars-surprisingly-galaxy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.