

Breaking the chain—catalyzing a green future for chemistry

November 6 2017



Fig.1: TEM image of CeO2-supported ruthenium nanoparticles catalyst. Yellow circles show Ru nanoparticles. Credit: Osaka University



Osaka University researchers create catalyst for refining chemicals in plant waste, allowing a green way to produce valuable raw materials.

The fight against climate change is a call-to-arms for industry. We currently rely on fossil fuels, a major source of the greenhouse gas CO2, not only for energy but also to create chemicals for manufacturing. To ween our economies off this dependency, we must find a new source of "green" <u>raw materials</u> so that factories and laboratories can run without producing and emitting CO2.

Now, a research team at Osaka University has discovered how to create valuable chemicals from clean sources. They used biomass, essentially waste from plant <u>materials</u>. Biomass is rich in organic molecules – long chains of carbon atoms attached to oxygen. Existing methods can break the carbon–oxygen bonds in these molecules to create, for example, raw materials for plastics. However, breaking the carbon–carbon bonds, in order to shorten the molecular chains, is harder; extreme temperatures are needed, and often yield unwanted products.

The method developed at Osaka is based on a <u>new catalyst</u>. Catalysts allow reactions to occur, without being consumed themselves. They are often based on metals, and the new example is no exception – it consists of atomically small particles of ruthenium, a metal related to iron, sitting on a material called cerium oxide.





Fig.2: Selective C-C bond cleavages of biogenic chemicals catalyzed by the CeO2-supported ruthenium nanoparticle catalyst. Credit: Osaka University

After creating their catalyst, the researchers tested it on levulinic acid (LA) from biomass. LA was reacted to break C-C bond at 150 degrees Celsius – hot for some, but mild by industrial standards. The reaction product was 2-butanol, an important <u>chemical</u> for manufacturing solvents. "This is the first time that 2-butanol has been made in this green way, using LA," study first author Tomoo Mizugaki explains. "Traditionally, it is made from butene, which comes from highly polluting oil refineries."

Buoyed by this, the team tested their catalyst on other biomass chemicals. A range of valuable products was obtained. Crucially, the reactions always broke carbon–<u>carbon</u> bonds. This allowed them to produce, for example, cyclohexanol, an important chemical in the



manufacture of nylon.



Fig.3: The new synthetic routes for valuable chemicals using the CeO2-supported ruthenium nanoparticle catalyst. Credit: Osaka University

X-ray and microscope studies confirmed that the combination of



ruthenium, <u>cerium oxide</u> and water was vital for the reaction to occur. The new <u>catalyst</u> therefore fills an important gap in the chemist's toolbox of reactions.

"We hope this method helps all sectors of industry obtain raw materials from non-fossil sources," corresponding author Kiyotomi Kaneda says. "We need a radical change in thinking, so that bio-derived chemicals are considered as primary options in manufacturing."

More information: Tomoo Mizugaki et al. New Routes for Refinery of Biogenic Platform Chemicals Catalyzed by Cerium Oxide-supported Ruthenium Nanoparticles in Water, *Scientific Reports* (2017). DOI: 10.1038/s41598-017-14373-1

Provided by Osaka University

Citation: Breaking the chain—catalyzing a green future for chemistry (2017, November 6) retrieved 6 August 2024 from <u>https://phys.org/news/2017-11-chaincatalyzing-green-future-chemistry.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.