


# Dawn explores Ceres' interior evolution

November 9 2017

---



This image made with data from NASA's Dawn spacecraft shows pit chains on dwarf planet Ceres called Samhain Catenae. Credit: NASA/JPL-Caltech/UCLA/MPS/DLR/IDA

Surface features on Ceres—the largest world between Mars and Jupiter—and its interior evolution have a closer relationship than one might think.

A recent study, published in *Geophysical Research Letters*, analyzed

Ceres' surface [features](#) to reveal clues about the dwarf planet's interior evolution. Specifically, the study explored linear features—the chains of pits and small, secondary craters common on Ceres.

The findings align with the idea that, hundreds of millions (up to a billion) years ago, materials beneath Ceres' surface pushed upward toward the exterior, creating [fractures](#) in the crust.

"As this material moved upward from underneath Ceres' surface, portions of Ceres' outer layer were pulled apart, forming the fractures," said Jennifer Scully, lead study author and associate of the Dawn science team at NASA's Jet Propulsion Laboratory in Pasadena, California.

The indication of upwelling material under Ceres' surface allows for another perspective in establishing how the [dwarf planet](#) may have evolved.

## **Searching for a Needle in a Haystack**

Dawn scientists generated a map of over 2,000 linear features on Ceres greater than 0.6 mile (one kilometer) in length that are located outside of impact craters. The scientists interpreted Dawn's observations of two kinds of linear features to further understand their connection to the upwelling material. Secondary [crater](#) chains, the most common of the linear features, are long strings of circular depressions created by fragments thrown out of large [impact craters](#) as they formed on Ceres. Pit chains, on the other hand, are surface expressions of subsurface fractures.

Among the two features, only pit chains provide insight into Ceres' interior evolution. Scully said the study's greatest challenge was differentiating between secondary crater chains and pit chains. Although the features are strikingly similar, researchers were able to distinguish

between them based on their detailed shapes. For example, secondary craters are comparatively rounder than pit chains, which are more irregular. In addition, pit chains lack raised rims, whereas there is usually a rim around secondary craters.

## How the Features Formed

While it is possible that the freezing of a global subsurface ocean formed the fractures, this scenario is unlikely, as the locations of pit chains are not evenly dispersed across Ceres' surface. It is also unlikely that the fractures formed by stresses from a large impact because there is no evidence on Ceres of impacts substantial enough to generate fractures of that scale. The most probable explanation, according to the Dawn scientists, is that a region of upwelling material formed the pit chains. The material may have flowed upward from Ceres' interior because it is less dense than surrounding materials.

Dawn scientists look forward to seeing how these characteristics will help other researchers model Ceres' interior evolution, which can test whether upwelling may have occurred near the fractures.

**More information:** J. E. C. Scully et al. Evidence for the Interior Evolution of Ceres from Geologic Analysis of Fractures, *Geophysical Research Letters* (2017). [DOI: 10.1002/2017GL075086](https://doi.org/10.1002/2017GL075086)

Provided by Jet Propulsion Laboratory

Citation: Dawn explores Ceres' interior evolution (2017, November 9) retrieved 26 April 2024 from <https://phys.org/news/2017-11-dawn-explores-ceres-interior-evolution.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.