

New discovery to accelerate development of salt-tolerant grapevines

November 23 2017



Credit: CC0 Public Domain

A recent discovery by Australian scientists is likely to improve the sustainability of the Australian wine sector and significantly accelerate the breeding of more robust salt-tolerant grapevines.



With funding from Wine Australia, a team of scientists from the ARC Centre of Excellence in Plant Energy Biology at the University of Adelaide and CSIRO Agriculture and Food identified genes expressed in grapevine roots that limit the amount of sodium – a key component of salt – that reaches berries and leaves. The research has been published this week in the journal *New Phytologist*.

'Berries that contain too much sodium may be unsuitable for wine production and this can lead to vineyards remaining unpicked, resulting in financial losses for vineyard owners,' says Dr Sam Henderson, co-first author of the study, from the University of Adelaide.

'We set out to determine why some grapevines accumulate salt and others don't, and found a specific mutation in a sodium transport protein found in grapevine roots, which prevents it from working effectively. This leads to more salt leaking into the shoots of vines from the soil,' Dr Henderson says.

While low levels of salt can improve the flavour of wine, in excess it can lead to unpalatable tastes, reduce fruit yield and damage the long-term health of grapevines – it is a problem experienced in premium wine regions around the world. In Australia's broader agriculture, food and wine sectors, issues caused by salinity have been estimated to cost in excess of \$1 billion each year.

'By comparing the DNA of different grapevines we identified a specific gene that is associated with sodium exclusion from shoots,' says co-first author Dr Jake Dunlevy from CSIRO.

'This discovery has allowed us to develop genetic markers that are being used to breed more salt-tolerant grapevine rootstocks, allowing new genotypes to be screened at the seedling stage rather than through lengthy and expensive field-based vineyard trials.'



'Traditionally, winegrape rootstocks have been developed in wine producing regions in the United States and Europe. This new research supports a breeding program to combine multiple beneficial traits in grapevines using conventional breeding, to develop robust rootstocks specifically for Australian conditions and support the local <u>wine</u> sector's sustainability well into the future,' says Dr Liz Waters, Wine Australia's General Manager Research, Development and Extension.

More information: Sam W. Henderson et al. Functional differences in transport properties of natural HKT1;1 variants influence shoot Na+ exclusion in grapevine rootstocks, *New Phytologist* (2017). <u>DOI:</u> <u>10.1111/nph.14888</u>

Provided by University of Adelaide

Citation: New discovery to accelerate development of salt-tolerant grapevines (2017, November 23) retrieved 2 May 2024 from <u>https://phys.org/news/2017-11-discovery-salt-tolerant-grapevines.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.