

# Feral animals pose major threat to Outback, climate change study finds

November 1 2017

---



Lead author Dr Aaron Greenville captures the splendour of central Australia's Simpson Desert. This photo was taken in the south-west of Queensland. Credit: Dr Aaron Greenville

A study of changing rainfall and bushfire patterns over 22 years has

found - in addition to a likely decrease in cover of the dominant plant spinifex - feral animals pose a major threat to seed-eating rodents.

Scientists at the University of Sydney have analysed up to 22 years of long-term monitoring data on plants and animals in central Australia to project how changing rainfall and wildfire patterns, because of climate change, will influence desert wildlife.

The research from the University of Sydney's School of Life and Environmental Sciences found there could be decreases in the cover of the dominant plant, spinifex, and a resulting decrease in seeding - and that introduced cats and foxes pose one of the major threats to seed-eating rodents.

Spinifex is a major habitat type across central Australia - from Queensland to Western Australia, including in the Simpson Desert where this study was conducted.

The findings are being published in the Royal Society's journal *Open Science*.

Some estimates have suggested that up to one in six species are at risk from [climate change](#). Predators can take advantage of more open habitats for hunting, which puts more pressure on wildlife.



An introduced cat in the study area. Credit: Desert Ecology Research Group, School of Life and Environmental Sciences, University of Sydney

Lead author Dr Aaron Greenville said their model suggested that if removing introduced cats and foxes, the rodent population would increase by almost one in 10.

"The dingo plays a complex positive role for wildlife in arid Australia," said Dr Greenville.



The desert mouse, *Pseudomys desertor*, is listed as Critically Endangered in NSW but is not threatened in Queensland or for the IUCN Red List. It is one of the species of rodents that researchers studied as part of their work. Credit: Dr Aaron Greenville

**More information:** Aaron C. Greenville et al. Desert mammal populations are limited by introduced predators rather than future climate change, *Royal Society Open Science* (2017). [DOI: 10.1098/rsos.170384](https://doi.org/10.1098/rsos.170384)

Provided by University of Sydney

Citation: Feral animals pose major threat to Outback, climate change study finds (2017, November 1) retrieved 28 June 2024 from <https://phys.org/news/2017-11-feral-animals-pose-major-threat.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.