


Gravitational waves could shed light on the origin of black holes

November 30 2017, by Kevin Stacey



The LIGO experiment has made several detections of colliding black holes. Future gravitational wave experiments might detect such events much further back in time, which could shed light on how black holes form. Credit: The SXS (Simulating eXtreme Spacetimes) Project

A new study published in *Physical Review Letters* outlines how scientists could use gravitational wave experiments to test the existence of primordial black holes, gravity wells formed just moments after the Big

Bang that some scientists have posited could be an explanation for dark matter.

"We know very well that black holes can be formed by the collapse of large stars, or as we have seen recently, the [merger](#) of two neutron stars," said Savvas Koushiappas, an associate professor of physics at Brown University and coauthor of the study with Avi Loeb from Harvard University. "But it's been hypothesized that there could be black holes that formed in the very early [universe](#) before stars existed at all. That's what we're addressing with this work."

The idea is that shortly after the Big Bang, quantum mechanical fluctuations led to the density distribution of matter that we observe today in the expanding universe. It's been suggested that some of those density fluctuations might have been large enough to result in black holes peppered throughout the universe. These so-called primordial black holes were first proposed in the early 1970s by Stephen Hawking and collaborators but have never been detected—it's still not clear if they exist at all.

The ability to detect gravitational waves, as demonstrated recently by the Laser Interferometer Gravitational-Wave Observatory (LIGO), has the potential to shed new light on the issue. Such experiments detect ripples in the fabric of spacetime associated with giant astronomical events like the collision of two black holes. LIGO has already detected several black hole mergers, and future experiments will be able to detect events that happened much further back in time.

"The idea is very simple," Koushiappas said. "With future gravitational wave experiments, we'll be able to look back to a time before the formation of the first stars. So if we see black hole merger events before stars existed, then we'll know that those black holes are not of stellar origin."

Cosmologists measure how far back in time an event occurred using redshift—the stretching of the wavelength of light associated with the expansion of the universe. Events further back in time are associated with larger redshifts. For this study, Koushiappas and Loeb calculated the redshift at which black hole mergers should no longer be detected assuming only stellar origin.

They show that at a redshift of 40, which equates to about 65 million years after the Big Bang, merger events should be detected at a rate of no more than one per year, assuming stellar origin. At redshifts greater than 40, events should disappear altogether.

"That's really the drop-dead point," Koushiappas said. "In reality, we expect merger events to stop well before that point, but a redshift of 40 or so is the absolute hardest bound or cutoff point."

A redshift of 40 should be within reach of several proposed gravitational wave experiments. And if they detect merger events beyond that, it means one of two things, Koushiappas and Loeb say: Either primordial black holes exist, or the early universe evolved in a way that's very different from the standard cosmological model. Either would be very important discoveries, the researchers say.

For example, primordial black holes fall into a category of entities known as MACHOs, or Massive Compact Halo Objects. Some scientists have proposed that dark matter—the unseen stuff that is thought to comprise most of the mass of the universe—may be made of MACHOs in the form of primordial black holes. A detection of primordial black holes would bolster that idea, while a non-detection would cast doubt upon it.

The only other possible explanation for [black hole mergers](#) at redshifts greater than 40 is that the universe is "non-Gaussian." In the standard

cosmological model, matter fluctuations in the early universe are described by a Gaussian probability distribution. A merger detection could mean matter fluctuations deviate from a Gaussian distribution.

"Evidence for non-Gaussianity would require new physics to explain the origin of these fluctuations, which would be a big deal," Loeb said.

The rate at which detections are made past a [redshift](#) of 40—if indeed such detections are made—should indicate whether they're a sign of [primordial black holes](#) or evidence for non-Gaussianity. But a non-detection would present a strong challenge to those ideas.

More information: Savvas M. Koushiappas et al, Maximum Redshift of Gravitational Wave Merger Events, *Physical Review Letters* (2017). [DOI: 10.1103/PhysRevLett.119.221104](https://doi.org/10.1103/PhysRevLett.119.221104)

Provided by Brown University

Citation: Gravitational waves could shed light on the origin of black holes (2017, November 30) retrieved 10 April 2024 from <https://phys.org/news/2017-11-gravitational-black-holes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
