

Study finds Texas's annual risk of extreme rainfall will rise from 1 to 18 percent

November 13 2017, by Jennifer Chu



“You’re rolling the dice every year,” says professor Kerry Emanuel. “And we believe the odds of a flood like Harvey are changing.” Pictured is an aerial view of Houston during the Hurricane Harvey flooding. Credit: Massachusetts Institute of Technology

As the city of Houston continues to recover and rebuild following the

historic flooding unleashed by Hurricane Harvey, the region will also have to prepare for a future in which storms of Harvey's magnitude are more likely to occur.

A new MIT study, published online this week in the *Proceedings of the National Academy of Sciences*, reports that as [climate change](#) progresses, the city of Houston, and Texas in general, will face an increasing risk of devastating, Harvey-scale rainfall.

According to the study, the state of Texas had a 1 percent chance of experiencing rainfall of Harvey's magnitude for any given year between 1981 and 2000. By the end of this century, the annual probability of Hurricane Harvey's record rainfall returning to Texas will rise to 18 percent, if the growth of [greenhouse gas emissions](#) to the atmosphere continues unmitigated.

If the risk for such an event during this century increased in a steady, linear fashion, it would mean that there was a 6 percent chance of having Harvey's magnitude of rainfall in Texas this year.

"You're rolling the dice every year," says study author Kerry Emanuel, the Cecil and Ida Green Professor of Atmospheric Science and co-director of the Lorenz Center at MIT. "And we believe the odds of a flood like Harvey are changing."

When the past isn't a guide

In the wake of a large disaster, Emanuel says it is natural, and in some cases essential, to ask whether and how soon such an event will occur again.

"Suppose you're the mayor of Houston, and you've just had a terrible disaster that cost you an unbelievable fortune, and you're going to try

over the next few years to put things back in order in your city," Emanuel says. "Should you be putting in a more advanced storm-sewer system that may cost billions of dollars, or not? The answer to that question depends upon whether you think Harvey was a one-off—very unlikely to happen any time in the next 100 years—or whether it may be more common than you thought."

Looking at historical records of extreme rainfall will not provide much insight into the future, Emanuel says. That's because past measurements have been spotty and difficult to extrapolate across larger regions, and the period over which rainfall data have been recorded is relatively short. What's more, climate change is shifting the odds in terms of the frequency of high-intensity storms around the world.

"If the underlying statistics are changing, the past may not be a good guide to the future," Emanuel notes in the paper.

Instead, scientists are turning to climate models to try and forecast the future of storms like Harvey. But there, challenges also arise, as models that simulate changing climate at a global scale do so at relatively coarse resolution, of around hundreds of kilometers, while hurricanes require resolutions of a few kilometers.

"[Climate models] do simulate slushy [hurricane](#)-like storms, but they're very poorly resolved," Emanuel says. "We don't have the computational firepower to resolve storms like hurricanes in today's climate models."

Hurricanes, embedded

Emanuel and his colleagues had previously devised a technique to simulate hurricane development in a changing climate, using a specialized computational [model](#) they developed that simulates hurricanes at high spatial resolutions. The model is designed so that they

can embed it within coarser [global climate](#) models—a combination that results in precise simulations of hurricanes in the context of a globally changing climate.

Emanuel used the team's technique to model past and future hurricane activity for both the city of Houston and the state of Texas. To do so, he first embedded the hurricane model in three gridded climate analyses—simulations of global climate, based on actual data from the past—to simulate hurricane activity near Houston between 1980 and 2016.

He randomly seeded each climate model with hundreds of thousands of "proto-hurricanes," or early-stage storms, the majority of which naturally peter out and don't grow to become full-fledged hurricanes. Of the remaining storms, he focused on the 3,700 storms that passed within 300 kilometers of Houston between 1980 and 2016. He then noted the frequency of storms that produced 500 millimeters of rainfall or more—the amount of rain that was initially estimated immediately following Hurricane Harvey.

During this historical period, he calculated that the probability of a Harvey-like storm producing at least 500 millimeters of rain in Houston was around once in 2,000 years. Such an event, he writes, was "'biblical' in the sense that it likely occurred around once since the Old Testament was written."

Stormy odds

To get a sense for how this probability, or risk of such a storm, will change in the future, he performed the same analysis, this time embedding the hurricane model within six global [climate models](#), and running each model from the years 2081 to 2100, under a future scenario in which the world's climate changes as a result of unmitigated

growth of greenhouse gas emissions.

While Houston's yearly risk of experiencing a 500-millimeter rainfall event was around 1 in 2,000 at the end of the last century, Emanuel found the city's annual odds will increase significantly, to one in 100 by the end of this century.

When he performed the same set of analyses for Texas as a whole, he found that, at the end of the 20th century, the state faced a 1 percent risk each year of experiencing a Harvey-scale [storm](#). By the end of this century, that annual risk will increase to 18 percent. If this increase happens linearly, he calculates that this year, the state's odds were at about 6 percent—a sixfold increase since the late 20th century.

"When you take a very, very rare, extreme [rainfall](#) event like Hurricane Harvey, and you shift the distribution of rain toward heavier amounts because of [climate](#) change, you get really big changes in the probability of those rare events," Emanuel says. "People have to understand that damage is usually caused by extreme events."

Emanuel hopes that the study's results will help city planners and government officials to decide where and how to rebuild and fortify infrastructure, as well as whether to recode building standards to stand up to stronger storms and more damaging floods.

"We're seeing for Texas an event whose annual probability was 1 percent at the end of the last century, and it might be 18 percent by the end of this century," Emanuel says. "That's a huge increase in the probability of that event. So, people had better plan for that."

More information: Assessing the present and future probability of Hurricane Harvey's rainfall, *PNAS*,
www.pnas.org/cgi/doi/10.1073/pnas.1716222114

Provided by Massachusetts Institute of Technology

Citation: Study finds Texas's annual risk of extreme rainfall will rise from 1 to 18 percent (2017, November 13) retrieved 27 April 2024 from <https://phys.org/news/2017-11-texas-annual-extreme-rainfall-percent.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.