


# Cellular division strategy shared across all domains of life

December 19 2017, by Leah Burrows

---



SEAS researchers have found that these pink-hued archaea — called *Halobacterium salinarum* — use the same mechanisms to maintain size as bacteria and eukaryotic life, indicating that cellular division strategy may be shared across all domains of life. Credit: Alexandre Bison/Harvard University

The three domains of life—archaea, bacteria, and eukarya—may have more in common than previously thought.

Over the past several years, Ariel Amir, Assistant Professor in Applied Mathematics at the Harvard John A. Paulson School of Engineering and Applied Sciences (SEAS) has been studying how cells regulate size. In previous research, he and his collaborators found that *E. coli* ([bacteria](#)) and budding yeast (eukaryote) use the same [cellular mechanisms](#) to ensure uniform cell sizes within a population.

Now, with a team of collaborators including Ethan Garner, the John L. Loeb Associate Professor of the Natural Sciences at Harvard, and Amy Schmid, Assistant Professor of biology at Duke University, Amir found that archaea use the very same mechanism.

The research is published in *Nature Microbiology*.

"These findings raise really interesting questions about how cellular mechanics evolved independently across all three domains of life," said Amir. "Our results will serve as a useful foundation for, ultimately, understanding the molecular mechanisms and evolution of [cell cycle](#) control."

Archaea are single-celled microorganisms that inhabit some of Earth's most extreme environments, such as volcanic hot springs, oil wells and salt lakes. They are notoriously difficult to cultivate in a lab and, as such, are relatively understudied.



Archaea inhabit some of Earth's most extreme environments, such as this salt lake in Bolivia. Credit: Ariel Amir/Havard SEAS

"Archaea are unique because they blend a lot of the characteristics of both bacteria and eukaryotes," said Dr. Yejin Eun, first author of the paper. "Archaea resemble [bacterial cells](#) in size and shape but their cell cycle events—such as division and DNA replication—are a hybrid between eukaryotes and bacteria."

The researchers studied *Halobacterium salinarum*, an extremophile that lives in high-salt environments. They found that like bacteria and budding yeast, *H. salinarum* controls its size by adding a constant volume between two events in the cell cycle. However, the researchers found that *H. salinarum* are not as precise as *E.coli* and there was more

variability in cell division and growth than in bacterial [cells](#).

"This research is the first to quantify the cellular mechanics of size regulation in [archaea](#)," said Amir. "This allows us to quantitatively explore how these mechanisms work, and build a model that explains the variability within the data and the correlations between key properties of the cell cycle. Eventually, we hope to understand just what makes this cellular [mechanism](#) so popular across all domains of life."

**More information:** Ye-Jin Eun et al. Archaeal cells share common size control with bacteria despite noisier growth and division, *Nature Microbiology* (2017). [DOI: 10.1038/s41564-017-0082-6](https://doi.org/10.1038/s41564-017-0082-6)

Provided by Harvard University

Citation: Cellular division strategy shared across all domains of life (2017, December 19) retrieved 27 June 2024 from <https://phys.org/news/2017-12-cellular-division-strategy-domains-life.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--