


Combination of warmer water, chemical exposure intensifies harmful effects in a coastal fish

December 15 2017, by Chris Branam



A drawing of an adult silverside. Credit: © Andrea Dingeldein

Warmer water temperatures, combined with exposure to chemicals already known to be harmful to aquatic life, could threaten organisms that have temperature-sensitive sex determination.

Researchers found that inland silversides (*Menidia beryllina*) exposed to endocrine disrupting compounds (EDCs) exhibited altered sex ratios, lower fertility rates and deformities. Further, the effects were more acute in the fish exposed at the warmer water temperatures predicted by the Intergovernmental Panel on Climate Change.

The study was published in the journal *Scientific Reports*.

"This may be a serious issue for this species and others," said Susanne Brander, an aquatic toxicologist at Oregon State University who co-authored the study. "We saw fewer offspring in multiple generations at the warmer temperatures across the board. With altered sex ratio and altered reproduction, there is certainly potential for population-level effects."

Inland silversides are small—adults are about 4 inches long—that are native to estuaries in eastern North America and the Gulf of Mexico and have been introduced into California. They primarily feed on zooplankton and are an important prey species for a variety of fish and birds.

The compounds studied by the researchers were the insecticide bifenthrin and ethinylestradiol (EE2), a synthetic estrogen found in almost all combined forms of [birth control pills](#).

A large portion of EE2 is not absorbed by the body and is excreted in urine. Wastewater treatment plants generally aren't equipped to eliminate such chemicals and they end up in rivers, and eventually estuaries. Bifenthrin is used to control insects in homes, orchards and nurseries.

In the study, three generations of silversides were exposed to a nanogram per liter of bifenthrin and EE2, in water at 22 degrees Celsius (71.6 degrees Fahrenheit) and 28 degrees Celsius (82.4 degrees Fahrenheit).

"The exposure levels were equivalent to a drop in an Olympic-sized swimming pool," Brander said. "Our scenario was fish exposed in a runoff event, for example in spring when people have just applied pesticide to their homes and there is a lot of rain."

Each generation was exposed to the chemicals for a 35-day period.

The change in [temperature](#) was important, Brander said, because silversides, as well as some other vertebrate species, have evolved to produce more [females](#) in colder water and more males in warmer [water](#), giving the females a longer time to grow and produce more eggs. But in the study, more females were produced in both chemical exposures at warmer temperatures, giving the females less time to grow. Smaller females tend to produce fewer eggs, Brander said.

There were also significant levels of tailfin deformities, with the effects being worse at warmer temperatures. A malformed tailfin makes it difficult for the fish to avoid predators.

More information: Bethany M. DeCourten et al. Combined effects of increased temperature and endocrine disrupting pollutants on sex determination, survival, and development across generations, *Scientific Reports* (2017). [DOI: 10.1038/s41598-017-09631-1](https://doi.org/10.1038/s41598-017-09631-1)

Provided by Oregon State University

Citation: Combination of warmer water, chemical exposure intensifies harmful effects in a coastal fish (2017, December 15) retrieved 2 May 2024 from <https://phys.org/news/2017-12-combination-warmer-chemical-exposure-effects.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.