

# How invasive weeds can make wildfires hotter and more frequent

December 21 2017, by Tim Curran, George Perry And Sarah Wyse

---



Mixed grill: Burning combinations of invasive and native plants helps us understand how invasive plants make fires hotter and more likely. Credit: Sarah Wyse, CC BY-ND

Over the past year the global media has been full of reports of catastrophic fires in [California](#), the Mediterranean, [Chile](#) and elsewhere. One suggested reason for increases in catastrophic wildfires has been [human-induced climate change](#). Higher temperatures, drier weather and windier conditions all increase the impact of fires.

While climate change indeed raises the risk of wildfires, our [research](#) shows that another way humans can change patterns of fire activity is by introducing flammable [plants](#) to new environments.

Plantations of highly flammable exotic [species](#), such as pines and eucalypts, probably helped to fuel the recent [catastrophic fires](#) in Portugal and in [Chile](#). In arid regions, such as parts of the [US southwest](#), the introduction of exotic grasses has transformed shrublands, as fires increase in severity.

## **Invasive plants and fire**

How do invasive plants change fire patterns? We burned species mixtures (aka "[mixed grills](#)") on our [plant barbecue](#) to help find out.

Invasive plants are responsible for [changing the patterns of fire activity](#) in many ecosystems around the world. In particular, invasive species can lead to [hotter](#) and [more frequent](#) fires.

Invasive plants can also [reduce fire frequency and fire intensity](#), but there are [fewer examples](#) of this occurring worldwide.



Samples of *Hakea sericea* (foreground) and *Kunzea robusta* (rear) arranged on the grill of our plant barbecue. Credit: Sarah Wyse, CC BY-ND

One of the main ways flammable invasive plants can have long-lasting impacts on an ecosystem comes from [positive fire-vegetation feedbacks](#). Such feedbacks can occur when a flammable weed invades a less fire-prone ecosystem. By [changing the available fuel](#) the invader makes fires more likely and often hotter.

If the invading species has [characteristics](#) that allow it to outcompete native species after a fire, then it will further dominate the ecosystem. Such traits include thick bark, the ability to resprout following fire, or

seeds that survive burning. This invasion will likely lead to more fires, changing the species composition and function of the ecosystem in a "fire begets fire" cycle. Extreme examples of this dynamic are where flammable [grasses](#) or [shrubs](#) invade forests, leading to loss of the forest ecosystems.

## **Mixed grills**

We wanted to understand how invasive plants interact with other species when burned in combination. To explore the mechanisms underpinning such feedbacks, we examined how invasive plants might [change the nature of a fire](#) when burned together with native species.

We collected 70cm shoots of four globally invasive species (of both high and low flammability) and burned them in pairwise combinations with New Zealand native trees and shrubs to determine which characteristics of a fire could be attributed to the [invasive plants](#).

We found that overall flammability was largely driven by the most flammable species in the mixture, showing how highly flammable weeds could set in motion fire-vegetation feedbacks.

We established that a greater difference in flammability between the two species led to a larger influence of the more flammable species on overall flammability. This outcome suggests weeds that are much more flammable than the invaded community can have larger impacts on fire patterns.

Importantly, we also showed the influence of the highly flammable species was independent of its biomass, meaning highly flammable weeds may impact community flammability even at low abundances.



Measuring how long a fire burns on our plant barbecue. Credit: Tom Etherington, CC BY-ND

When we looked closer at the different components of flammability (combustibility, ignitability, consumability and sustainability) we found some important nuances in our results.

While the maximum temperature reached in our burns (combustibility) and the ignition speed (ignitability) were both most influenced by the more flammable species, consumability (the amount of biomass burned) and sustainability (how long the fire burns) were equally influenced by both the more flammable and less flammable species.

In short, more flammable weeds will cause a fire to ignite more quickly and burn hotter.

However, less flammable species can reduce the duration of a fire compared to when a more flammable species is burnt alone. These results could have important ecological implications, as the [longer a fire burns the more likely it is to kill plants](#): low-flammability plants could reduce this impact.

## Managing weeds to reduce fire impacts

Even low abundances of highly flammable invasive weeds could set in motion positive fire-vegetation feedbacks that lead to drastic changes to ecosystems. If this result holds when our shoot-scale experiments are repeated using field trials, then land managers should work quickly to remove even small infestations of highly flammable species, such as gorse (*Ulex europaeus*) and prickly hakea (*Hakea sericea*).

Conversely, the role of low flammability plants in extinguishing fires further supports the suggestion that the strategic planting of such species across the landscape as "green firebreaks" could be a useful [fire](#) management tool.

In any case, our "mixed grill" study further highlights the role of exotic plants in fuelling hotter wildfires.

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by The Conversation

Citation: How invasive weeds can make wildfires hotter and more frequent (2017, December 21)

retrieved 2 May 2024 from

<https://phys.org/news/2017-12-invasive-weeds-wildfires-hotter-frequent.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.