

## Scientists synthesize a new phosphor

December 22 2017



Crystal structure of the compound. Credit: Yana Baklanova (co-author of the research study)

Chemists and physicists from Ural Federal University and Institute of Chemistry of Solids of the Ural Branch of Russian Academy of Sciences



have synthesized a new compound that may be used to convert UV radiation into visible light and change laser radiation range. The work was published in the *Journal of Luminescence*.

The article authors worked with lithium-containing compounds with garnet structure. The chemical formula of such compositions is  $Li_xLa_3M_2O_{12}$ , where M is zirconium, niobium, tin, tellurium, hafnium, or tantalum and x equals three to seven. A number of publications about materials based on these systems have been published lately. However, their optical properties remained understudied.

From previous publications, chemists knew that many hafnium compounds had photo-, X-ray-, and radioluminescence. Thus, ceriumdoped hafnium garnet  $Ca_3Hf_2SiAlO_{12}$  phosphor shows a broad cyan emission under 400 nm excitation (on the border between visible and UV radiation). The authors of the work decided to synthesize lithiumcontaining cerium-doped hafnium garnet (Li<sub>7</sub>La<sub>3</sub>Hf<sub>2</sub>O<sub>12</sub>:Ce<sub>3</sub><sup>+</sup>) to obtain one more composition that emits in the visible spectrum.

After synthesizing the necessary <u>compounds</u>, the chemists studied its properties. Using X-ray diffraction, they determined the compound's structural characteristics. They studied its microstructure using scanning electron microscopy. To determine the relationship between reflection, excitation and luminescence spectra on the dopant concentration and temperature, the chemists carried out a complex analysis of optical spectroscopy. They found a correlation between separate emission bands and F+ centers.

"After detailed analysis of all obtained characteristics and using EPR spectroscopy, we managed to prove that the increase of cerium concentration led to the reduction in the intensity of photoluminescence associated with intrinsic luminescence centers in Li7La3Hf2O12 represented by excitons (quasiparticles as bound states of electron and



hole) in regular crystal lattice nods (Hf-O)," said Yana Baklanova, a coauthor of the article, and senior scientific associate of Institute of Chemistry of Solids of Ural Branch of Russian Academy of Sciences.

The compound, with other dopants, may be used in the manufacture of medical and optic devices and monitoring systems,

"A  $\text{Li}_7\text{La}_3\text{Hf}_2\text{O}_{12}$  compound doped with europium and neodymum or holmium (Eu<sub>3</sub><sup>+</sup> and Nd<sub>3</sub><sup>+</sup>/Ho<sub>3</sub><sup>+</sup>) may by used to convert UV radiation into the visible spectral range and monochromatic laser radiation into the infrared range with shortwave- and middle-infrared spectral range. IR <u>radiation</u> is widely used in medicine, various optical systems, and also for air pollution analysis," concluded Alexey Ishchenko, a co-author of the article, assistant professor and senior scientific associate of Ural Federal University.

Provided by Ural Federal University

Citation: Scientists synthesize a new phosphor (2017, December 22) retrieved 1 May 2024 from <u>https://phys.org/news/2017-12-scientists-phosphor.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.