

Strongly anisotropic spin relaxation observed in graphene

December 8 2017



Credit: ICN2

Researchers of the ICN2 Physics and Engineering of Nanodevices Group, led by ICREA Prof. Sergio O. Valenzuela, have unambiguously demonstrated the anisotropic nature of spin relaxation in graphene when



interfaced with transition metal dichalcogenides (TMDC). The paper, titled "Strongly anisotropic spin relaxation in graphene–transition metal dichalcogenide heterostructures at room temperature," was published this week in *Nature Physics* with lead author L. Antonio Benítez.

Spin is the <u>intrinsic angular momentum</u> of subatomic particles. Although with no real equivalent in classical physics, it can be used much like charge to store, manipulate and transport information. Graphene is known to transport <u>electron spins</u> very effectively over large distances. However, manipulating these spins is made difficult by the lack of any means of controlling them externally. The spin-orbit coupling (SOC) interaction offers just such a means, though in graphene, this must be 'borrowed' from other materials via the proximity effect.

Using an experimental approach developed over the last two years by the ICN2 group, the researchers observed that spins behave differently upon reaching the graphene/TMDC bilayer depending on their orientation. Inplane spins were found to be very sensitive to the proximity-induced SOC, relaxing and losing their orientation as much as 10 times faster than out-of-plane spins. This strongly anisotropic spin relaxation was further observed to be a consequence of the spin-valley coupling also imprinted onto the graphene from the TMDC.

Crucially, all observations were carried out at <u>room temperature</u>, making them of direct relevance to future technological applications.

These results suggest that a graphene/TMDC system could be used as a spin filter, allowing the detection of small orientation changes. They also represent the first step to achieving external control over the propagation of spins in graphene, and offer an interesting starting point for the exploration of coupled spin-valley phenomena and spin-valleytronics device concepts.



More information: L. Antonio Benítez et al. Strongly anisotropic spin relaxation in graphene–transition metal dichalcogenide heterostructures at room temperature, *Nature Physics* (2017). <u>DOI:</u> <u>10.1038/s41567-017-0019-2</u>

Provided by ICN2

Citation: Strongly anisotropic spin relaxation observed in graphene (2017, December 8) retrieved 3 May 2024 from <u>https://phys.org/news/2017-12-strongly-anisotropic-graphene.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.