


Researchers find way to give advanced notice for hailstorms

January 30 2018, by Tom Parisi



New research finds a strong relationship between a wavy jet-stream pattern over the Pacific Ocean and severe U.S. hailstorms. Credit: Victor Gensini, NIU

A new study identifies a method for predicting the likelihood of damaging hailstorms in the United States—up to three weeks in advance.

Hail is the most economically destructive hazard posed by severe thunderstorms, producing on average billions of dollars in U.S. losses each year, including damage to roofs, homes and especially crops.

"We found a really strong relationship between jet stream patterns over the Pacific Ocean and U.S. hail frequency," said Victor Gensini, a meteorologist at Northern Illinois University in DeKalb, Illinois and lead author of the new study. "In simple terms, when the jet stream is really wavy, the likelihood of experiencing hail greatly increases."

The new study was accepted for publication in *Geophysical Research Letters*, a journal of the American Geophysical Union.

Two years ago, Gensini led research on a method to predict the likelihood of U.S. tornado activity weeks in advance. Last year, of 26 long-range (two- to three-week) forecasts for increased, average or below average U.S. tornado activity, more than half were "spot on," Gensini said. Most of the other predictions were only slightly off.

The new study is an extension of the tornado research, suggesting a similar method can be used in sub-seasonal forecasts of hailstorms. "There's a high degree of correlation between environments that produce hail and tornadoes, but not all storms produce both hazards," Gensini said.

While the method would be used to forecast hail activity for the country in general, portions of Texas, Oklahoma, Arkansas, Kentucky, Missouri, Mississippi, Tennessee, Illinois and Indiana are most vulnerable to the phenomenon.

Gensini and Allen examined hail observations from national storm data for the period of 1979 to 2016. They compared those events with changes in the Global Wind Oscillation (GWO) index, a collection of

climate and weather information that measures atmospheric angular momentum, or the degree of waviness in the jet stream.

The GWO index has eight distinct phases. Four of those phases were reliable predictors of increased inland hail activity during peak storm seasons, according to the study.

"There is a strong connection between the GWO and U.S. hail frequency," said John Allen, a meteorologist at Central Michigan University in Mt. Pleasant, Michigan and co-author of the new study. "This relationship helps to understand what is driving hail variability, and explains to a large degree when we are likely to experience active and inactive periods during the spring and fall.

During the summer, however, the GWO is unreliable as other smaller-scale meteorological processes tend to dominate local weather conditions.

"We will be testing the relationships this spring when severe weather season ramps up," Gensini said.

"We're starting to demonstrate more clearly a pathway to increase the lead time for severe weather forecasts, now with both hailstorms and tornadoes," he added. "We keep adding cinderblocks to the methodology, and it's slowly becoming robust."

More information: Vittorio A. Gensini et al. United States Hail Frequency and the Global Wind Oscillation, *Geophysical Research Letters* (2018). [DOI: 10.1002/2017GL076822](https://doi.org/10.1002/2017GL076822)

Provided by Northern Illinois University

Citation: Researchers find way to give advanced notice for hailstorms (2018, January 30)
retrieved 3 May 2024 from <https://phys.org/news/2018-01-advanced-hailstorms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.